

# Novel Role for the *yceGH* Tellurite Resistance Genes in the Pathogenesis of *Bacillus anthracis*

Sarah E. Franks,<sup>a</sup> Celia Ebrahimi,<sup>b</sup> Andrew Hollands,<sup>b</sup> Cheryl Y. Okumura,<sup>b</sup> Raffi V. Aroian,<sup>c</sup> Victor Nizet,<sup>b,d</sup> Shauna M. McGillivray<sup>a</sup>

Department of Biology, Texas Christian University, Fort Worth, Texas, USA<sup>a</sup>; Department of Pediatrics,<sup>b</sup> Division of Biological Sciences,<sup>c</sup> and Skaggs School of Pharmacy and Pharmaceutical Sciences,<sup>d</sup> University of California San Diego, La Jolla, California, USA

*Bacillus anthracis*, the causative agent of anthrax, relies on multiple virulence factors to subvert the host immune defense. Using *Caenorhabditis elegans* as an infection model, we screened approximately 5,000 transposon mutants of *B. anthracis* Sterne for decreased virulence. One of the attenuated mutants resulted in loss of expression of *yceG* and *yceH*, the last two genes in a six-gene cluster of tellurite resistance genes. We generated an analogous insertional mutant to confirm the phenotype and characterize the role of *yceGH* in resistance to host defenses. Loss of *yceGH* rendered the mutants more sensitive to tellurite toxicity as well as to host defenses such as reactive oxygen species and the cathelicidin family of antimicrobial peptides. Additionally, we see decreased survival in mammalian models of infection, including human whole blood and in mice. We identify a novel role for the *yceGH* genes in *B. anthracis* Sterne virulence and suggest that *C. elegans* is a useful infection model to study anthrax pathogenesis.

*Bacillus anthracis* is a Gram-positive, spore-forming bacterium and the etiological agent of the deadly disease anthrax. The most lethal form of the disease, inhalational anthrax, occurs when inhaled spores are phagocytosed by resident macrophages or dendritic cells and are carried to regional lymph nodes (1). En route, the bacteria germinate, escape the phagolysosome, and break out of the cell to replicate extracellularly, eventually leading to massive septicemia, shock, and death (2). The hardy nature of the spores, their relative ease of dissemination, and the high mortality rate make *B. anthracis* a paramount bioterrorism concern.

*B. anthracis* must avoid an array of host bacterial defenses during the course of infection. Major plasmid-encoded virulence factors include the lethal and edema toxins and a poly-D-glutamate capsule (2). Although the anthrax toxins and capsule play central roles in disease pathogenesis, increasing evidence indicates that chromosomal genes also contribute to virulence. Sequencing of the *B. anthracis* genome identified a number of genes that share homologies to known virulence factors of other pathogens (3). Subsequently, studies have coupled targeted mutagenesis with rodent infection models to identify chromosomal virulence genes, including the *dltABCD* operon for lipoteichoic acid modification (4), the *asbAB* siderophore (5), the ABC transporter *mntA* (6), the nitric oxide synthase *nos* (7), the ClpXP protease (8), the purine biosynthesis gene *purH* (9), and the stress resistance gene *hrtA* (10).

Forward genetic screens using transposon-based mutagenesis systems can be a powerful method to identify novel bacterial virulence genes. We successfully employed this system in our identification of the *clpX* gene, encoding part of the intracellular protease ClpXP, as necessary for *B. anthracis* virulence (8). One challenge of this approach is designing a productive screen. Due to ethical and logistical constraints, it is not possible to perform large-scale virulence screens in vertebrate models. Instead, surrogate *in vitro* phenotypes that directly or indirectly correlate with virulence are often examined, for example, loss of hemolysis in identification of *clpX* (8).

A more amenable, whole-animal model to screen for virulence factors emerged with the finding that *B. anthracis* can infect and

kill the nematode *Caenorhabditis elegans* (11). *C. elegans* has gained attention as a whole-animal infection model in recent years (12) because it possesses an evolutionarily conserved innate immune system with host defenses such as antimicrobial peptides (AMPs) (13) and reactive oxygen species (ROS) (14), as well as amenable traits, including small size, rapid generation time, and production of large numbers of genetically identical offspring. *C. elegans* has served as a host model system for several bacterial pathogens, including *Pseudomonas aeruginosa* and *Staphylococcus aureus*, with mutants subsequently confirmed to have decreased virulence in murine challenge experiments (12).

In the present study, we employ a transposon mutant library that we developed previously (8) to screen for *B. anthracis* Sterne mutants unable to infect *C. elegans*. Our most highly attenuated mutant had a transposon insertion in the second-to-last gene of a six-gene cluster of bacterial tellurite resistance genes. We find that these last two genes are important not only for tellurite resistance but also for resistance to specific reactive oxygen species (ROS) such as hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>), the cathelicidin family of antimicrobial peptides, and human whole-blood killing. This increased susceptibility correlates with reduced virulence of our transposon mutant in *C. elegans* and murine models of infection.

## MATERIALS AND METHODS

**Bacterial strains and culture conditions.** *B. anthracis* Sterne (pXO1<sup>+</sup> pXO2<sup>-</sup>) was grown in brain heart infusion (BHI) medium (Hardy Diagnostics) at 37°C with continuous shaking. We used antibiotic selection at the following concentrations: 50 µg/ml kanamycin (Kan) and 5 µg/ml of

Received 16 December 2013 Accepted 18 December 2013

Published ahead of print 23 December 2013

Editor: S. R. Blanke

Address correspondence to Shauna M. McGillivray, s.mcgillivray@tcu.edu.

Copyright © 2014, American Society for Microbiology. All Rights Reserved.

doi:10.1128/IAI.01614-13

erythromycin (Em). Construction of the transposon mutant library was described previously (8).

**Construction of targeted insertional mutant and complementation plasmid.** Approximately 600 bp of *yceG* sequence was amplified using the forward primer 5'-ATGCAAGCTTATGTCGAATCCCTTGGCGTT-3' and the reverse primer 5'-GTCAGAATTCCGACGATCTGGAAATGGC T-3', which contained 5' extensions with restriction sites for HindIII and EcoRI, respectively. This amplicon was then cloned into the temperature-sensitive plasmid pHY304 (15), a derivative of pVE6007 (16), which replicates in a temperature-sensitive manner. Transformants were grown initially at 30°C under Em selection and then shifted to the nonpermissive temperature of 37°C to force plasmid integration into the bacterial chromosome. Integration was confirmed by PCR using the pHY304-specific primer 5'-ACGACTCACTATAGGGCGAATTGG-3' and the primer 5'-CCTTCTGGTAATCGCTTATGTGGC-3', which is located downstream of the original amplicon.

The *yceG* and *yceH* genes were amplified as one amplicon using the forward primer 5'-ATGGTACCGGACAAGCTAAAGAAGAAGTTGCA GCT-3' and the reverse primer 5'-ATGAGCTCTTAGTCTTCTCCCTTC TTATCGGTGTT-3' and cloned into the pDCerm expression vector (17) at the KpnI and SacI restriction sites. This plasmid was designated pGH and transformed into the TN1 strain.

**C. elegans infection assay.** The *C. elegans* *glp-4(bn2)* strain, which is sterile at 25°C, was maintained using standard techniques (18). The *C. elegans* infection assay previously described (11) was modified to accommodate a high-throughput 96-well assay. Two days prior to the infection assay, a synchronous population of L1 worms was seeded onto nematode growth plates and fed *Escherichia coli* strain OP50. The worms were incubated at 25°C for 48 h to sterilize them and allow them to reach the young adult stage. The worms were then collected from plates, washed twice, and resuspended in 1 ml of S medium (19). Five microliters of this medium (containing approximately 5 to 10 worms) was then transferred to 190 µl of S medium containing Kan and 10 µg/ml Cry5B. Five microliters of an overnight *B. anthracis* culture grown in 96-well plates was then added for a final volume of 200 µl. The plates were wrapped in Parafilm and incubated at 28°C. Wells were assessed at 48 and 72 h. Nematodes were counted as infected only if they were filled with bacteria, which can be detected as worms being dark, rigid, and immobile. Wells with more than 60% of worms surviving were noted for further confirmation in a larger volume (24-well plates) in an independent assay.

**Identification of transposon insertions.** Transposon insertions were identified using two different methods. DNA flanking the transposon insertion site was identified by single-primer PCR as described previously (8). Alternatively, the insertion site was identified using a Y-linker method (20). The transposon insertion site was confirmed using the transposon-specific primer 5'-TATGCATTTAATACTAGCGACG-3' and one of the following primers located downstream of the insertion site: TN1, 5'-GT GAAGCGAAAGGAATCGAAGT-3'; TN2, 5'-TGTACTTGAACATCAG TCGATTCTCCA-3'; TN3, 5'-TACTGCCAATTATGAATAAAC CA-3'; TN4, 5'-CATGCGTGGCAAGTTA-3'; TN5, 5'-TCTTAGTGCAG GGAAACATAACTGA-3'; TN6, 5'-TCCCACCATAATTCCATGACA CC-3'; TN7, 5'-CAGAGATTACAATAGGTTCGGCT-3'; TN8, 5'-TTCG TCTTATCAGGATTACTCAGTTAG-3'; TN9, 5'-TGTCATCGGAAGC ATAATCCT-3'; TN10, 5'-CAAGAAATGCTCGTTGGCTATC-3'; and TN11, 5'-TCTTTGCATCGGAACATTGCT-3'.

**RT-PCR.** Sixteen-hour bacterial cultures were lysed using Lysing Matrix B beads (MP Biomedical) by being pulsed two times for 45 s at 6 m/s in a bead beater (MP Biomedical). RNA was purified using the RNeasy kit (Qiagen) followed by DNase treatment using the Turbo DNA-free kit (Life Technologies). For reverse transcription-PCR (RT-PCR) shown in Fig. 2B, RNA was reverse transcribed using Superscript III reverse transcriptase (Life Technologies) at 55°C with one of the following gene-specific primers: RT 1, 5'-CTCGCTACTTCCACACTTGC-3'; RT 2, 5'-GA ATACGCCAGCGATACCGC-3'; or RT 3, 5'-CCTCAACTGCTTGCCTA ACTT-3'. PCR was performed at 95°C for 30 s, 54°C for 30 s, and 72°C for

1 min for 30 cycles with the following primer combinations: A (fwd, 5'-ACTATCGCCTGAGAGGCCAC-3', and rev, 5'-GCTCCTCCAATAC CAGTTCCA-3'), B (fwd, 5'-TAGAGGGCATTCATATAGTGCTGG-3', and rev, 5'-GATCCCAACCTAACGCCAACCTG-3'), C (fwd, 5'-GTGAA GGCGCTGGTGATGAC-3', and rev, 5'-GAAATCCTCTGCACCTGAA ACT-3'), D (fwd, 5'-TACGATGGAGAAGGACGCAG-3', and rev, 5'-C CTGTCGGTTATCTCCTGT-3'), E (fwd, 5'-AGGTGGTTAGGTGC GCTTG-3', and rev, 5'-CATGAGCTCCTCTTCTGC-3'), F (fwd, 5'-GCGGTATCGCTGGCGTATTC-3', and rev, 5'-CATTAGGAACGCC AAGATTGCG-3'), and G (fwd, 5'-TTACAACATGGGATTGAGAGG-3', and rev, 5'-CAATTTCACGGGCCATCGTT-3'). For the RT-PCR shown in Fig. 2C, RNA was reverse transcribed using a high-capacity cDNA reverse transcription kit (Life Technologies). PCR was performed at 95°C for 30 s, 54°C for 30 s, and 72°C for 1 min and limited to 26 cycles to maintain semiquantitative conditions. The following primers were used: *yceC* Fwd, 5'-CAATCAGGTGGGTTCTGT-3'; *yceC* Rev, 5'-TGA TGCCATATACAATTCCCTCTGT-3'; *yceG* Fwd, 5'-CGTCCACAAT ATGACCTGACAG-3'; *yceG* Rev, 5'-GAGCTACAGGTGTATAGGAAC G-3'; *yceH* Fwd, 5'-TGGACCGATTGATAGCAA-3'; *yceH* Rev, 5'-CGA CTTTCTTCACGCTCACGC-3'; *fusA* Fwd, 5'-AAGCTGGTGGTGTG AAGCAC-3'; *fusA* Rev, 5'-TTCCAATCAGCTCTCCTGAAG-3'. All RT-PCRs were repeated with at least 3 different RNA preparations; a representative image is shown.


**Growth curves.** Growth curves were started using equal amounts of early-log-phase parental and mutant bacterial strains, and bacteria were grown at 37°C with shaking in BHI with the indicated amount of potassium tellurite (Sigma). Absorbance was recorded at 600 nm. Growth curves were repeated at least three times; a representative graph is shown.

**Disk diffusion assays.** Parental or mutant *B. anthracis* stationary-phase cultures (16 h) were swabbed onto BHI agar plates. A 6-mm sterile paper disk (Becton, Dickinson) containing a specific antimicrobial was placed on the agar, the plate was incubated overnight at 37°C, the zone of inhibition was measured, and the area was calculated. Tellurite disks were prepared by soaking the disk in 10 mg/ml of potassium tellurite. Antibiotic disks containing 30 µg chloramphenicol, 30 µg tetracycline, and 10 U penicillin were obtained from Becton, Dickinson. For antibiotic disk diffusion performed with tellurite, the agar plates contained 30 µg/ml of potassium tellurite.

**Oxidant assays.** For the growth assays, *B. anthracis* parental and mutant strains were grown to early log phase; diluted 1:2 (H<sub>2</sub>O<sub>2</sub>), 1:10 (hypochlorite), or 1:100 (methyl viologen); and incubated with the indicated amounts of each specific oxidant (Sigma). The final optical density (OD) after overnight incubation (≈16 h) was used as the measure of growth. A MIC assay was performed with 0, 10, and 40 mM methyl viologen (Sigma). For the H<sub>2</sub>O<sub>2</sub> killing assay, bacteria were grown to an optical density of 0.4 at 600 nm and concentrated 10× in phosphate-buffered saline (PBS) and the concentrated bacteria were then diluted 1:2 into 0.1% H<sub>2</sub>O<sub>2</sub> in PBS for a final concentration of 0.05% H<sub>2</sub>O<sub>2</sub>. Bacteria were incubated at 37°C. At the indicated time, an aliquot was removed and diluted 1:10 in PBS containing 3,000 U catalase (Sigma) to quench residual H<sub>2</sub>O<sub>2</sub> and surviving bacteria were enumerated by serial dilution plating.

**Antimicrobial peptide assays.** *B. anthracis* parental and mutant strains were grown to early log phase, diluted 1:10 in RPMI plus 5% Luria-Bertani broth, and incubated with 1.6 µM LL-37 (Anaspec) overnight at 37°C under static conditions. MIC assays were also performed using RPMI plus 5% Luria-Bertani broth and log-phase bacteria diluted to 1:20 with either 0, 125, and 250 µg/ml nisin; 0, 250, 500, and 1,000 µg/ml bacitracin; or 0, 25, 50, and 100 µg/ml polymyxin (all from Sigma) or log-phase bacteria diluted to 1:100 and 0, 0.4, and 0.8 µM HNP-2 (Anaspec).

**Whole-blood assay.** Blood collected from three different healthy donors (use and procedures approved by the University of California, San Diego, Human Research Protections Program) was incubated with 10<sup>5</sup> CFU *B. anthracis* in a total volume of 500 µl and rotated at 37°C. After 15



**FIG 1** Transposon mutants have attenuated virulence in a *C. elegans* model of infection. *C. elegans* mortality rate after 72 h of infection with WT *B. anthracis* Sterne or transposon mutants (TN1 to -11). Data from at least three independent experiments are combined and presented as means  $\pm$  standard errors of the means. \*,  $P < 0.05$ ; \*\*,  $P < 0.01$ ; \*\*\*,  $P < 0.001$ , from WT by one-way analysis of variance followed by Tukey's *post hoc* analysis.

min, aliquots were removed, blood was lysed in water, and surviving bacteria were enumerated by serial dilution plating.

**In vivo infection.** Bacteria were grown to an optical density (OD) of 0.4 (600 nm), washed in PBS, and resuspended at a 1:30 dilution in PBS (approximately  $7 \times 10^5$  CFU/ml). Parental (wild-type [WT]) and mutant (TN1) *B. anthracis* Sterne strains were mixed at a 1:1 ratio, and 0.1 ml was injected intravenously via the lateral tail vein into 8- to 10-week-old female CD1 mice. After 2 days, animals were euthanized and the kidneys were isolated and weighed. Kidneys were homogenized (twice for 1 min each at 6,000 rpm) using 1-mm zirconia-silica beads (BioSpec Products) in 2-ml screw-cap tubes containing sterile PBS using a MagNA lyser (Roche). Surviving bacteria were enumerated by serial dilution plating on BHI (total CFU) and BHI Kan50 plates (TN1 CFU). WT CFU was determined by subtracting the Kan-resistant CFU from the total CFU, and bacterial counts were calculated as CFU/g kidney.

**Bioinformatics and statistics.** Protein sequences were aligned using the ClustalW program found at [http://npsa-pbil.ibcp.fr/cgi-bin/npsa\\_automat.pl?page=/NPSA/npsa\\_clustalw.html](http://npsa-pbil.ibcp.fr/cgi-bin/npsa_automat.pl?page=/NPSA/npsa_clustalw.html). The following sequences were used: the *B. anthracis* *yceCDEFGH* operon gene symbol BAS0385 through BAS0390, the R478 Ter operon with GenBank accession number **U59239.2**, and the *Listeria monocytogenes* TelA gene symbol lmo1967. Statistical analysis was performed using GraphPad Prism software.

## RESULTS

### Identification of an attenuated *B. anthracis* transposon mutant.

We have recently demonstrated that under the proper assay conditions, including supplementation with an exogenous pore-forming toxin, *B. anthracis* Sterne can infect *C. elegans* (11). Distinguishing between *B. anthracis*-infected and uninfected worms is based on visual appearance. Infected nematodes are rigid and dark and lack internal structures as they are filled with *B. anthracis* spores (11). Conversely, healthy animals are motile and curvilinear, displaying typical discernible internal structures. We screened approximately 5,000 *B. anthracis* transposon mutants created previously (8) for their ability to establish an infection in *C. elegans* using a high-throughput 96-well screening format. This is a non-saturating screen; however, in our previous use of this library, we found that 5,000 mutants is a sufficient number to identify promising targets (8). Under our infection conditions, the mortality rate of *C. elegans* with wild-type (WT) *B. anthracis* Sterne was approximately 75%. We identified 11 transposon mutants that yielded a *C. elegans* mortality rate of 40% or less (Fig. 1) and determined the site of disruption (Table 1).


The most consistently attenuated mutant, designated TN1, induced a *C. elegans* mortality rate of less than 20% and contains a disruption in a putative tellurite resistance gene (BAS0389). Tellurite ( $\text{TeO}_3^{2-}$ ) is a water-soluble form of tellurium and is highly toxic to bacteria. Tellurite resistance genes are found in a number of bacterial species, although they have not been linked to pathogenesis (21). TN2 has a disruption in a gene containing BNR repeats (also known as ASP-box motifs) that are commonly found in glycosyl hydrolases such as sialidases, which have been linked to bacterial pathogenesis (22). TN3, TN4, and TN5 are all inserted in intergenic regions, and the disrupted gene is assumed to be the gene downstream of the insertion site. TN4 disrupts a predicted protease production regulatory protein, which could have implications in virulence. TN5 is in the middle of a gene cluster of conserved hypothetical proteins with little information on potential function. TN3 and TN7 also disrupt conserved hypothetical proteins of unknown function. These conserved hypothetical pro-

**TABLE 1** Summary of attenuated mutants identified in *C. elegans* infection screen

| Mutant | Disrupted gene <sup>a</sup>            | Gene symbol | Sequence surrounding insertion site <sup>b</sup>   |
|--------|--|-------------|--|
| TN1    | Putative tellurite resistance gene     | BAS0389     | ACGTATCATTTGGATGGTGACAGGAAAGTCGTATATTCTTATTTCTTATTGCTGG<br>TTGTGATGTGCTTATTATCACCAGAAAGGA            |
| TN2    | BNR repeat domain protein              | BAS2814     | AACCAAATCTATCAGTCGCATTGTAAGAAATTGAATATCATTATTCAAATATCACC<br>TCATTCTTATCATATATATTCTGATGCTT            |
| TN3    | Conserved hypothetical protein         | BAS0642     | GCACITGAAAAGATAAAACACAAAGTGAAAATTCTGTTCACCTAAATAATAGAATAGTAAAAG<br>TAGCTCTTCTAGAAATAGAAGGGCTACT      |
| TN4    | Protease production regulatory protein | BAS0976     | TGCATAAAAACACGGCATATCTATGTGATATGCCGTGTCAAGGTCATTTATATAAAAATTTGG<br>ATATGCAAAAACAAGCAGGAATTGTTAC      |
| TN5    | Hypothetical protein                   | BAS0432     | ATTGAAAATTGGCAGCATCTGATCCAGACATTATATGTAAGTAAAAATAAGGGATGG<br>AAATCATGAGAGAACGTGAAAATTGGGATGT         |
| TN6    | Spore germination protein              | BAS3369     | GTATTAATAGGACCATATGACCTTTCATCTTCAACCGATAGATGATTACAGCTCAGACCAAT<br>GATTCCATCTTATACGTTACTCGTTCA        |
| TN7    | Conserved hypothetical protein         | BAS0645     | TAGAGATAGGAGATTCTATCTGTTGCCATAGTAAAGCTGGAGAATTATGTGAGCGGTTCC<br>TTATATCGTGTGAAAGAAGAGAGATTGTTGG      |
| TN8    | Sensor histidine kinase                | BAS0869     | GAGAGAAAAGAATGTTCTCTTGTGACGGATATTGGCAGCATATTGTATTGAGTAATAGAT<br>AGAGGGATAGGAATACCGCAGAAAAATAAC       |
| TN9    | PTS, IIB component                     | BAS2290     | GTGATCATCCCGCCGAAGCATATACGCCACTTGGCGGACCGACTTTATTAAAGACTGTAATGA<br>TTAATCGTGTAGGGGAACCGTTATGAATAGAAG |
| TN10   | Acetyl-CoA hydrolase                   | BAS1726     | ATTATTGAGAATTGTGACATCTAATGTATCTGATCAACTACGAGCTTATTGAGGAAGCGAAA<br>CAAGAGGTGGACAAACTCCTCATATTAGAGA    |
| TN11   | Oxidoreductase                         | BAS0712     | ATATTAGTAAATATGTCGCCAGCAATATCCGCAGCAAGGACTAGAGTATTACAGCAGAAC<br>AGTTAGAAAAGACTTTCGTTAATATTTC         |

<sup>a</sup> Abbreviations: PTS, phosphotransferase system; CoA, coenzyme A.

<sup>b</sup> Insertion site is shown in bold.



**FIG 2** Organization of the *yce* tellurite resistance gene cluster in *B. anthracis*. (A) Organization of the tellurite resistance gene cluster in *B. anthracis* with number of amino acids (aa) listed below each gene. The upward arrow indicates the site of transposon and plasmid insertion for the transposon mutant 1 and the insertional mutant 1 (TN1/IM1). Leftward arrows indicate primers used for reverse transcription (RT 1, RT 2, or RT 3). Arrowheads show location of PCR pairs A to G. The bent arrow indicates the predicted promoter region. (B) PCR using cDNA from indicated RT primers. Letters correlate with primer combinations shown in panel A. (C) Semiquantitative RT-PCR expression of the *yceC*, *yceG*, and *yceH* genes from cDNA prepared from WT, TN1, IM1, or TN1 complemented with an expression plasmid containing *yceG* and *yceH* (pGH).

teins are on either side of phospholipase C and sphingomyelinase, which have been implicated in *B. anthracis* virulence (23). While it is unlikely that either TN3 or TN7 directly disrupts these virulence genes, it is possible that the conserved hypothetical proteins have a functional cooperation with phospholipase C and/or sphingomyelinase that would account for this coincidence. TN6 inserts in an apparent operon containing spore germination proteins. TN9, TN10, and TN11 all disrupt metabolic genes and may impair general bacterial fitness. TN8 is found in a sensor histidine kinase (BAS0869) and likely also disrupts the response regulator (BAS0870) and lipoprotein (BAS0871) located directly downstream in a putative operon. Notably, after a short intergenic region of 220 bp, the next gene in this region is another putative tellurite resistance protein (BAS0872). We tried to assess whether TN1 and TN8 had similar phenotypes; however, this was complicated by a growth defect in TN8 (data not shown) that may itself account for the decreased virulence seen in the *C. elegans* model.

Although several of the disrupted genes are potentially of interest and would be interesting to pursue in future studies, we have focused this study on TN1. Not only is TN1 highly attenuated in our *C. elegans* model, it also disrupts an intriguing gene family that has been speculated, although to our knowledge never proven, to have a role in bacterial virulence (21).

**Tellurite resistance genes in *B. anthracis*.** The *B. anthracis* genome contains a six-gene cluster of putative tellurite resistance genes (BAS3085 to BAS3090) in which the second-to-last gene is disrupted by the transposon insertion in TN1 (Fig. 2A, arrow). Homology with the corresponding proteins in *Bacillus subtilis* is high (Table 2), and we have maintained the gene nomenclature, *yceCDEFGH*, originally designated for *B. subtilis* (24). We used an RT-PCR approach for characterization of multigene operons (25) to test whether all 6 putative tellurite resistance genes were regulated within the same operon. Reverse transcription was performed using primer RT 1, RT 2, or RT 3 followed by PCR with

primer sets A to G, whose forward and reverse primers are located on neighboring genes (Fig. 2A). PCR amplification of the cDNA should occur only with genes expressed on the same transcript, although all primers amplified using genomic DNA (data not shown). As a negative control, we looked at expression between BAS0383 and BA3084, which are orientated in opposite directions and cannot be transcribed together. As expected, no amplification was seen using primer set A with cDNA from RT 2 (Fig. 2B) or cDNA from RT 3 (data not shown). We find that *yceCDEF* are transcribed together on the same operon; however, there is little to no expression using a primer set amplifying between *yceF* and *yceG* (Fig. 2B). Primers amplifying between *yceG* and *yceH* have strong expression, indicating that these genes are transcribed together. A very faint band is consistently seen between the upstream gene BAS0384 and *yceC*, although it is difficult to visualize. Therefore, while there may be some transcription of *yceCDEF* from the promoter upstream of BAS0384, the majority of expression is likely driven by a regulatory element between BAS0384 and *yceC*. We conclude that the putative tellurite resistance genes contained in *yceCDEF* and *yceGH* are regulated independently of one another.

At least five unique bacterial tellurite resistance ( $Te^r$ ) determi-

**TABLE 2** Homology of *B. anthracis* Yce proteins to other tellurite resistance proteins


| <i>B. anthracis</i> protein | <i>B. subtilis</i> protein(s) (% homology) | Ter/Tel protein(s) (% homology)       |
|-----------------------------|--|---------------------------------------|
| YceC                        | YceC (62.3)                                | TerE (40.1), TerD (37.0), TerZ (37.0) |
| YceD                        | YceD (68.6), YceE (67.0)                   | TerD (54.2), TerE (53.6)              |
| YceE                        | YceE (71.1), YceD (63.6)                   | TerD (55.9), TerE (52.6)              |
| YceF                        | YceF (64.3)                                | TerC (22.9)                           |
| YceG                        | YceG (41.4)                                |                                       |
| YceH                        | YceH (58.1)                                | TelA (24.1)                           |

nants have been identified (21). The *B. anthracis* YceCDEF and YceGH proteins have homology with two of them, TerZABCDEF and TelA (Table 2). The *terZABCDEF* operon was originally identified on the R478 plasmid, and homologues have been found in the chromosomes of numerous species, including *Yersinia pestis*, *E. coli*, *Bacillus subtilis*, and *Proteus mirabilis* (21, 24, 26, 27). The *telA* gene has been recently characterized in the Gram-positive pathogen *L. monocytogenes* (28). At the amino acid level, *B. anthracis* YceCDEF has homology to several Ter proteins (Table 2). YceG, the site of the transposon insertion, has little identity to any other tellurite resistance proteins, while YceH has 24.1% identity to the TelA protein found in *L. monocytogenes*.

To confirm the involvement of the *yceG* gene in the observed phenotype of the transposon mutant, we constructed an independent mutation (IM1) in approximately the same location using targeted insertional plasmid mutagenesis. RT-PCR confirmed that *yceG* gene expression was absent in both the TN1 and IM1 mutants relative to the wild type (Fig. 2C), whereas the first gene, *yceC*, is expressed in all three strains as expected. The last gene, *yceH*, is absent or substantially reduced in TN1 but present in IM1 (Fig. 2C). This differential regulation of *yceH* by TN1 and IM1 was also confirmed by quantitative PCR (qPCR) (data not shown). Therefore, IM1 functions as a genetic knockout of *yceG* whereas the transposon mutant functions as a double knockout of *yceG* and *yceH*. For complementation analysis, *yceG* and *yceH* were cloned on an expression plasmid (designated pGH) and transformed into the TN1 strain. Although transcription of the *yceG* and *yceH* genes was confirmed by RT-PCR in our complemented strain (Fig. 2C, lanes marked pGH), there was no change in phenotype from TN1 lacking the complementation plasmid in any subsequent assays (data not shown). This lack of complementation may result from improper stoichiometry of the proteins involved. No difference in expression of an unrelated control gene, *fusA*, was seen in any strains (data not shown).

Since *yceG* and *yceH* are homologous to putative tellurite resistance genes, we first tested susceptibility to potassium tellurite. We observed that both WT and mutant strains of *B. anthracis* Sterne form the characteristic black precipitate caused by reduction of tellurite ( $\text{TeO}_3^{2-}$ ) to tellurium ( $\text{Te}^0$ ) (Fig. 3A). However, there is a small but statistically significant increase in the zone of growth inhibition surrounding a potassium tellurite-impregnated disk in the TN1 and IM1 strains in comparison to WT (Fig. 3A). Exposure to potassium tellurite in liquid culture also results in slower growth for both TN1 and IM1 relative to the WT bacteria (Fig. 3B), although no difference in growth is seen in BHI alone (data not shown). Although other genes likely contribute, *yceGH* appears to play a minor role in tellurite resistance in *B. anthracis* Sterne.


***yceG* and *yceH* genes contribute to innate immune defense mechanisms.** Because increased susceptibility to tellurite toxicity cannot by itself explain the attenuated virulence of TN1 in *C. elegans*, we next investigated whether *yceGH* could play a role in defense against aspects of the host innate immune system. A critical innate immune defense of *C. elegans* is the production of ROS in response to pathogens (14). In mammalian systems, professional phagocytes such as macrophages and neutrophils generate ROS (29). Tellurite is also a strong oxidant (30) and increases intracellular ROS production in exposed bacteria (31, 32). Therefore, we determined whether *yceG* and *yceH* contribute to ROS resistance. Early-log-phase bacteria were incubated overnight in



**FIG 3** Loss of *yceG* leads to decreased tellurite resistance. Growth inhibition of WT *B. anthracis* Sterne (black bars), transposon mutant (TN1, white bars), or insertional mutant (IM1, gray bars) spread on plates and exposed to a tellurite-impregnated disk. The area of growth inhibition was calculated and presented as mean  $\pm$  standard deviation (right). \*,  $P < 0.05$ , or \*\*,  $P < 0.01$ , from WT by one-way analysis of variance followed by Tukey's *post hoc* analysis. The experiment was repeated at least three times, and representative pictures are shown (left). (B) Representative graph from a 24-hour growth curve in liquid BHI containing 30  $\mu\text{g}/\text{ml}$  of potassium tellurite.

BHI medium containing 0% and 0.02%  $\text{H}_2\text{O}_2$ . Although no difference in growth as measured by the final optical density was observed in BHI alone, addition of  $\text{H}_2\text{O}_2$  significantly attenuated growth for both the TN1 and IM1 mutants (Fig. 4A). This killing occurred relatively rapidly, as exposure to 0.05%  $\text{H}_2\text{O}_2$  killed all TN1 mutants by 45 min and reduced survival of IM1 by 100,000-fold in comparison to WT (Fig. 4B). While loss of *yceG* alone (IM1) appears sufficient to increase ROS sensitivity, *yceH* also likely contributes, since the phenotype seen in the TN1 mutant (loss of both *yceG* and *yceH*) is often stronger (compare survivals of TN1 and IM1 mutants in Fig. 4B). Additionally, we tested susceptibility for two additional ROS, sodium hypochlorite, the active ingredient in bleach, and superoxide generated by methyl viologen. As with  $\text{H}_2\text{O}_2$ , growth of TN1 and IM1 was significantly decreased compared with WT when bacteria were exposed to hypochlorite (Fig. 4C). However, no difference in survival was seen with superoxide (data not shown).

Antimicrobial peptides (AMPs) are a second innate immune defense mechanism employed by *C. elegans* (13). In *L. monocytogenes*, the tellurite resistance gene *telA* was identified in a transposon-based screen for susceptibility to the AMP nisin (28). Since *yceH* has homology to *telA* and is absent in our TN1 mutant, we tested whether TN1 and IM1 were more susceptible to several antimicrobial peptides, including the human cathelicidin LL-37, the  $\alpha$ -defensin HNP-2, nisin, bacitracin, and polymyxin. A signif-



**FIG 4** YceG is necessary for resistance to reactive oxygen species and human cathelicidin. (A, C, and D) Growth of WT *B. anthracis* Sterne (black bars), transposon mutant (TN1, white bars), or insertional mutant (IM1, gray bars) for 16 h in  $\text{H}_2\text{O}_2$  (A), hypochlorite (C), or cathelicidin (LL-37) (D). \*,  $P < 0.05$ , \*\*,  $P < 0.01$ , or \*\*\*,  $P < 0.001$ , from WT by one-way analysis of variance followed by Tukey's *post hoc* analysis. Data from at least three independent experiments are combined and presented as means  $\pm$  standard errors of the means. (B) Survival of WT *B. anthracis* Sterne (black circles), the transposon mutant (TN1, white circles), or the insertional mutant (IM1, gray circles) after incubation with 0.05%  $\text{H}_2\text{O}_2$  for 45 min. The experiment was repeated at least three times, and data from one representative experiment are presented as means  $\pm$  standard deviations.


ificant difference in survival was seen between WT and TN1 in LL-37 susceptibility (Fig. 4D); however, IM1 exhibited an intermediate phenotype (Fig. 4D). This is likely due to the fact that the TN1 insertion disrupts both *yceG* and *yceH* whereas IM1 disrupts only *yceG*. No difference in growth between WT *B. anthracis* Sterne, TN1, and IM1 exposed to HNP-2, nisin, bacitracin, or polymyxin was observed (data not shown). Thus, it appears that both *yceG* and *yceH* contribute to resistance to human cathelicidin; however, this did not represent a global change in membrane integrity as no difference in susceptibility was seen with any other AMP tested.

**Tellurite enhances susceptibility to antibiotics independently of *yceG* and *yceH*.** It has recently been reported that exposure to tellurite enhances susceptibility of *E. coli* to a broad range of antibiotics, including chloramphenicol, tetracycline, and ampicillin (33). Similarly, we observed a significantly increased zone of inhibition for *B. anthracis* Sterne in antibiotic disk diffusion assays for chloramphenicol, tetracycline, and penicillin when 30  $\mu\text{g}/\text{ml}$  of potassium tellurite was added to the agar plates in comparison to plates without potassium tellurite (Fig. 5A). We next asked whether mutations in the tellurite resistance genes would further enhance this sensitivity but found no difference in growth inhibition between the WT and TN1 mutant bacteria with any antibiotic in the absence (data not shown) or presence (Fig. 5B) of potassium tellurite. Therefore, while tellurite does increase *B. anthracis* susceptibility to antibiotics, this effect appears to be independent of *yceG* and *yceH*.

***yceG* and *yceH* contribute to *B. anthracis* pathogenicity in mammalian models of infection.** Since our transposon mutant is more susceptible to specific host defenses, we next assessed whether it was attenuated in mammalian models of infection. We found decreased survival of TN1 in human whole blood *ex vivo* (Fig. 6A), which contains a multitude of host defenses, including phagocytic cells, complement, and AMPs. We next tested whether TN1 was attenuated in an *in vivo* infection model. WT *B. anthracis* Sterne and the TN1 mutant were mixed at a 1:1 ratio and injected intravenously into mice. Bacterial survival in the kidneys was assessed when mice were moribund. Although equivalent numbers of WT and TN1 bacteria were injected (data not shown), survival of the transposon mutant was decreased about 10-fold in comparison to the WT bacteria (Fig. 6B), indicating that loss of *yceG* and *yceH* leads to attenuated virulence in a murine model of infection. Thus, *yceG* and *yceH* contribute to the overall virulence of *B. anthracis* in mammalian infection models. Furthermore, these data corroborate our results in the *C. elegans* screen (Fig. 1) and validate *C. elegans* as a screening model to identify *B. anthracis* Sterne mutants attenuated for virulence.


## DISCUSSION

Novel virulence genes in several pathogens have been identified through screens utilizing *C. elegans*, including, as we demonstrate here, *B. anthracis*. Our ability to successfully use *C. elegans* to identify virulence factors was verified by the identification of the transposon mutant TN1, which is more susceptible to ROS and



the human cathelicidin LL-37. TN1 has a disruption in *yceG*, the second-to-last gene in a six-gene cluster of tellurite resistance genes, that results in loss of expression of both *yceG* and *yceH*. Although originally identified as a putative operon (24), our results indicate independent regulation of *yceCDEF* and *yceGH*. There are at least five unique bacterial Te<sup>r</sup> determinants that have been identified, including the Ter proteins (similar to YceCDEF) and TelA (similar to YceH) (21). YceG remains poorly characterized with little sequence homology to other known tellurite resistance proteins. YceG has strong hydrophobic regions that may allow for interaction with the membrane, and across bacterial species, it is often found in association with TelA, indicating that these two proteins might be functionally linked (34). Despite their being widespread, surprisingly little is known regarding the actual function of any of the Te<sup>r</sup> genes.

As their name implies, the Te<sup>r</sup> genes are important for tellurite resistance in a number of bacterial species, including *E. coli*, *P. mirabilis*, and *Y. pestis* (21, 26, 27). While highly toxic to bacteria, tellurite is not necessarily commonly encountered, which raises the question of why these genes exist in such a wide range of species. It has been argued that the Te<sup>r</sup> genes are part of a general stress response system rather than solely tellurite resistance (21, 30, 34). In support of this, the Ter genes in *E. coli* are also implicated in resistance to bacteriophage and colicin (35). In *Y. pestis*, TerE and TerD proteins are upregulated along with other general



stress response proteins during intracellular replication in macrophages (26).

Our data show that loss of *yceG* and *yceH* in *B. anthracis* results in a small growth impairment in the presence of tellurite. However, the more pronounced phenotype is the increased susceptibility to host defenses such as ROS. Both our transposon and insertional mutants demonstrated increased sensitivity to H<sub>2</sub>O<sub>2</sub> and hypochlorite, the active ingredient in bleach. In a separate study in *B. anthracis*, *yceCDEF* were identified in an analysis of genes upregulated in response to H<sub>2</sub>O<sub>2</sub> (36). The Ter operon of *Proteus mirabilis* is also induced by H<sub>2</sub>O<sub>2</sub> and superoxide (27). Tellurite is also a strong oxidant (30), and tellurite toxicity has been linked to ROS production (31, 32). A recent comparative genomics study has found that the Ter genes are functionally linked to many enzymes involved in DNA processing and repair (34). It is therefore possible that these genes may play a role in repairing the DNA damage that would occur after exposure to oxidative stressors such as ROS or tellurite.

*B. anthracis* *yceG* and/or *yceH* is also important for resistance to the human cathelicidin LL-37. YceH has homology to TelA, which has been linked to resistance to nisin and some cell wall-targeting antibiotics in *L. monocytogenes* (28). Surprisingly, no difference in susceptibility was observed in the TN1 or IM1 mutants with any other AMP tested, including nisin, HNP-2, bacitracin, or polymyxin. This could be due to differences in mechanisms of action of the AMPs and/or in the responses of YceG/H. Although most AMPs form transmembrane pores, peptide insertion into the membrane can differ and some AMPs may have additional bactericidal mechanisms that could alter the bacterial response (37). In

*L. monocytogenes*, loss of TelA increases susceptibility to some cell wall-acting antibiotics (the cephalosporins, cefotaxime, and cefuroxime) but not all (no difference was seen with methicillin or oxacillin) (28). We also saw no contribution of the *yceG* or *yceH* gene to resistance to several common antibiotics, even when antibiotic susceptibility was magnified by coincubation with tellurite. Therefore, although *yceGH* may function as part of a general stress response system, there is still specificity in their mechanism of action.

A potential role of Te<sup>r</sup> genes in host defense has long been speculated (21), but to our knowledge, this is the first report directly linking any of them to increased susceptibility in mammalian infection models. We find that loss of *yceG* and *yceH* results in decreased survival of *B. anthracis* Sterne in human whole blood and a murine *in vivo* competition assay, both of which expose the bacteria to a synergistic array of host defenses. Taken together, our results and those of others support the argument that tellurite resistance genes are likely part of a more general chemical stress response system. Loss of these genes could render bacteria more susceptible to a variety of potential environmental stresses, including, but not limited to, exposure to strong oxidants. This would help explain their widespread and ubiquitous nature in a wide variety of bacterial species, including several human pathogens. Further study will clarify the roles and mechanism of action of this intriguing and mysterious family of genes.

## ACKNOWLEDGMENTS

This research was supported by NIH grant AI065993 (R.V.A. and V.N.). S.M.M. and C.Y.O. were supported by an IRCADA postdoctoral fellowship from NIGMS (GM068524), and S.M.M. was also supported by The Hartwell Foundation and through the Texas Christian University Research and Creative Activities Fund (60670).

## REFERENCES

1. Tournier JN, Paccani SR, Quesnel-Hellmann A, Baldari CT. 2009. Anthrax toxins: a weapon to systematically dismantle the host immune defenses. *Mol. Aspects Med.* 30:456–466. <http://dx.doi.org/10.1016/j.mam.2009.06.002>.
2. Dixon TC. 1999. Anthrax. *N. Engl. J. Med.* 341:815–826. <http://dx.doi.org/10.1056/NEJM199909093411107>.
3. Read TD, Peterson SN, Tourasse N, Baillie LW, Paulsen IT, Nelson KE, Tettelin H, Fouts DE, Eisen JA, Gill SR, Holtzapple EK, Okstad OA, Helgason E, Rilstone J, Wu M, Kolonay JF, Beanan MJ, Dodson RJ, Brinkac LM, Gwinn M, DeBoy RT, Madpu R, Daugherty SC, Durkin AS, Haft DH, Nelson WC, Peterson JD, Pop M, Khouri HM, Radune D, Benton JL, Mahamoud Y, Jiang L, Hance IR, Weidman JF, Berry KJ, Plaut RD, Wolf AM, Watkins KL, Nierman WC, Hazen A, Cline R, Redmond C, Thwaite JE, White O, Salzberg SL, Thomason B, Friedlander AM, Koehler TM, Hanna PC, Kolstø AB, Fraser CM. 2003. The genome sequence of *Bacillus anthracis* Ames and comparison to closely related bacteria. *Nature* 423:81–86. <http://dx.doi.org/10.1038/nature01586>.
4. Fisher N, Shetron-Rama L, Herring-Palmer A, Heffernan B, Bergman N, Hanna P. 2006. The dltABCD operon of *Bacillus anthracis* Sterne is required for virulence and resistance to peptide, enzymatic, and cellular mediators of innate immunity. *J. Bacteriol.* 188:1301–1309. <http://dx.doi.org/10.1128/JB.188.4.1301-1309.2006>.
5. Cendrowski S, MacArthur W, Hanna P. 2004. *Bacillus anthracis* requires siderophore biosynthesis for growth in macrophages and mouse virulence. *Mol. Microbiol.* 51:407–417. <http://dx.doi.org/10.1046/j.1365-2958.2003.03861.x>.
6. Gat O, Mendelson I, Chitlaru T, Ariel N, Altboum Z, Levy H, Weiss S, Grosfeld H, Cohen S, Shafferman A. 2005. The solute-binding component of a putative Mn(II) ABC transporter (MntA) is a novel *Bacillus anthracis* virulence determinant. *Mol. Microbiol.* 58:533–551. <http://dx.doi.org/10.1111/j.1365-2958.2005.04848.x>.
7. Shatalin K, Gusarov I, Avetissova E, Shatalina Y, McQuade LE, Lippard SJ, Nudler E. 2008. *Bacillus anthracis*-derived nitric oxide is essential for pathogen virulence and survival in macrophages. *Proc. Natl. Acad. Sci. U. S. A.* 105:1009–1013. <http://dx.doi.org/10.1073/pnas.0710950105>.
8. McGillivray SM, Ebrahimi CM, Fisher N, Sabet M, Zhang DX, Chen Y, Haste NM, Aroian RV, Gallo RL, Guiney DG, Friedlander AM, Koehler TM, Nizet V. 2009. ClpX contributes to innate defense peptide resistance and virulence phenotypes of *Bacillus anthracis*. *J. Innate Immun.* 1:494–506. <http://dx.doi.org/10.1159/000225955>.
9. Jenkins A, Cote C, Twenhafel N, Merkel T, Bozue J, Welkos S. 2011. Role of purine biosynthesis in *Bacillus anthracis* pathogenesis and virulence. *Infect. Immun.* 79:153–166. <http://dx.doi.org/10.1128/IAI.00925-10>.
10. Chitlaru T, Zaide G, Ehrlich S, Inbar I, Cohen O, Shafferman A. 2011. HtrA is a major virulence determinant of *Bacillus anthracis*. *Mol. Microbiol.* 81:1542–1559. <http://dx.doi.org/10.1111/j.1365-2958.2011.07790.x>.
11. Kho MF, Bellier A, Balasubramani V, Hu Y, Hsu W, Nielsen-LeRoux C, McGillivray SM, Nizet V, Aroian RV. 2011. The pore-forming protein Cry5B elicits the pathogenicity of *Bacillus* sp. against *Caenorhabditis elegans*. *PLoS One* 6:e29122. <http://dx.doi.org/10.1371/journal.pone.0029122>.
12. Sifri CD, Begun J, Ausubel FM. 2005. The worm has turned—microbial virulence modeled in *Caenorhabditis elegans*. *Trends Microbiol.* 13:119–127. <http://dx.doi.org/10.1016/j.tim.2005.01.003>.
13. Kato Y, Aizawa T, Hoshino H, Kawano K, Nitta K, Zhang H. 2002. abf-1 and abf-2, ASABF-type antimicrobial peptide genes in *Caenorhabditis elegans*. *Biochem. J.* 361:221–230. <http://dx.doi.org/10.1042/0264-6021:3610221>.
14. Chávez V, Mohri-Shiomi A, Garsin DA. 2009. Ce-Duox1/BLI-3 generates reactive oxygen species as a protective innate immune mechanism in *Caenorhabditis elegans*. *Infect. Immun.* 77:4983–4989. <http://dx.doi.org/10.1128/IAI.00627-09>.
15. Chaffin DO, Beres SB, Yim HH, Rubens CE. 2000. The serotype of type Ia and III group B streptococci is determined by the polymerase gene within the polycistronic capsule operon. *J. Bacteriol.* 182:4466–4477. <http://dx.doi.org/10.1128/JB.182.16.4466-4477.2000>.
16. Maguin E, Duwat P, Hege T, Ehrlich D, Gruss A. 1992. New thermo-sensitive plasmid for gram-positive bacteria. *J. Bacteriol.* 174:5633–5638.
17. Jeng A, Sakota V, Li Z, Datta V, Beall B, Nizet V. 2003. Molecular genetic analysis of a group A *Streptococcus* operon encoding serum opacity factor and a novel fibronectin-binding protein, SfbX. *J. Bacteriol.* 185:1208–1217. <http://dx.doi.org/10.1128/JB.185.4.1208-1217.2003>.
18. Brenner S. 1974. The genetics of *Caenorhabditis elegans*. *Genetics* 77:71–94.
19. Sulston JE, Brenner S. 1974. The DNA of *Caenorhabditis elegans*. *Genetics* 77:95–104.
20. Kwon YM, Ricke SC. 2000. Efficient amplification of multiple transposon-flanking sequences. *J. Microbiol. Methods* 41:195–199. [http://dx.doi.org/10.1016/S0167-7012\(00\)00159-7](http://dx.doi.org/10.1016/S0167-7012(00)00159-7).
21. Taylor DE. 1999. Bacterial tellurite resistance. *Trends Microbiol.* 7:111–115. [http://dx.doi.org/10.1016/S0966-842X\(99\)01454-7](http://dx.doi.org/10.1016/S0966-842X(99)01454-7).
22. Quistgaard EM, Thirup SS. 2009. Sequence and structural analysis of the Asp-box motif and Asp-box beta-propellers; a widespread propeller-type characteristic of the Vps10 domain family and several glycoside hydrolase families. *BMC Struct. Biol.* 9:46. <http://dx.doi.org/10.1186/1472-6807-9-46>.
23. Heffernan BJ, Thomason B, Herring-Palmer A, Shaughnessy L, McDonald R, Fisher N, Huffnagle GB, Hanna P. 2006. *Bacillus anthracis* phospholipases C facilitate macrophage-associated growth and contribute to virulence in a murine model of inhalation anthrax. *Infect. Immun.* 74:3756–3764. <http://dx.doi.org/10.1128/IAI.00307-06>.
24. Kumano M, Tamakoshi A, Yamane K. 1997. A 32 kb nucleotide sequence from the region of the lincomycin-resistance gene (22 degrees–25 degrees) of the *Bacillus subtilis* chromosome and identification of the site of the lin-2 mutation. *Microbiology* 143:2775–2782. <http://dx.doi.org/10.1099/00221287-143-8-2775>.
25. Gupta A. 1999. RT-PCR: characterization of long multi-gene operons and multiple transcript gene clusters in bacteria. *Biotechniques* 27:960–972.
26. Ponnusamy D, Hartson SD, Clinkenbeard KD. 2011. Intracellular *Yersinia pestis* expresses general stress response and tellurite resistance proteins in mouse macrophages. *Vet. Microbiol.* 150:146–151. <http://dx.doi.org/10.1016/j.vetmic.2010.12.025>.
27. Toptcheva A, Sisson G, Bryden LJ, Taylor DE, Hoffman PS. 2003. An inducible tellurite-resistance operon in *Proteus mirabilis*. *Microbiology* 149:1285–1295. <http://dx.doi.org/10.1099/mic.0.25981-0>.
28. Collins B, Joyce S, Hill C, Cotter PD, Ross RP. 2010. TelA contributes to the innate resistance of *Listeria monocytogenes* to nisin and other cell wall-acting antibiotics. *Antimicrob. Agents Chemother.* 54:4658–4663. <http://dx.doi.org/10.1128/AAC.00290-10>.

29. Lambeth JD. 2004. NOX enzymes and the biology of reactive oxygen. *Nat. Rev. Immunol.* 4:181–189. <http://dx.doi.org/10.1038/nri1312>.

30. Chasteen TG, Fuentes DE, Tantaleán JC, Vásquez CC. 2009. Tellurite: history, oxidative stress, and molecular mechanisms of resistance. *FEMS Microbiol. Rev.* 33:820–832. <http://dx.doi.org/10.1111/j.1574-6976.2009.00177.x>.

31. Pérez JM, Calderón IL, Arenas FA, Fuentes DE, Pradenas GA, Fuentes EL, Sandoval JM, Castro ME, Elías AO, Vásquez CC. 2007. Bacterial toxicity of potassium tellurite: unveiling an ancient enigma. *PLoS One* 2:e211. <http://dx.doi.org/10.1371/journal.pone.0000211>.

32. Tremaroli V, Fedi S, Zannoni D. 2007. Evidence for a tellurite-dependent generation of reactive oxygen species and absence of a tellurite-mediated adaptive response to oxidative stress in cells of *Pseudomonas pseudoalcaligenes* KF707. *Arch. Microbiol.* 187:127–135. <http://dx.doi.org/10.1007/s00203-006-0179-4>.

33. Molina-Quiroz RC, Muñoz-Villagrán CM, de la Torre E, Tantaleán JC, Vásquez CC, Pérez-Donoso JM. 2012. Enhancing the antibiotic antibacterial effect by sub lethal tellurite concentrations: tellurite and cefotaxime act synergistically in *Escherichia coli*. *PLoS One* 7:e35452. <http://dx.doi.org/10.1371/journal.pone.0035452>.

34. Anantharaman V, Iyer LM, Aravind L. 2012. Ter-dependent stress response systems: novel pathways related to metal sensing, production of a nucleoside-like metabolite, and DNA-processing. *Mol. Biosyst.* 8:3142–3165. <http://dx.doi.org/10.1039/c2mb25239b>.

35. Whelan KF, Colleran E, Taylor DE. 1995. Phage inhibition, colicin resistance, and tellurite resistance are encoded by a single cluster of genes on the IncH1I2 plasmid R478. *J. Bacteriol.* 177:5016–5027.

36. Pohl S, Tu WY, Aldridge PD, Gillespie C, Hahne H, Mäder U, Read TD, Harwood CR. 2011. Combined proteomic and transcriptomic analysis of the response of *Bacillus anthracis* to oxidative stress. *Proteomics* 11:3036–3055. <http://dx.doi.org/10.1002/pmic.201100085>.

37. Brogden KA. 2005. Antimicrobial peptides: pore formers or metabolic inhibitors in bacteria? *Nat. Rev. Microbiol.* 3:238–250. <http://dx.doi.org/10.1038/nrmicro1098>.