


Immunity

Normalizing Microbiota-Induced Retinoic Acid Deficiency Stimulates Protective CD8⁺ T Cell-Mediated Immunity in Colorectal Cancer

Graphical Abstract



Authors

Nupur Bhattacharya, Robert Yuan, Tyler R. Prestwood, ..., Daniel A. Winer, Joseph L. Napoli, Edgar G. Engleman

Correspondence

nupur26@stanford.edu (N.B.), edgareng@stanford.edu (E.G.E.)

In Brief

All-*trans*-retinoic acid (atRA) plays crucial roles in shaping intestinal immunity. Bhattacharya et al. show that, in the context of colon cancer, microbiota-induced intestinal inflammation alters atRA metabolism, leading to a colonic atRA deficit and exacerbation of colon carcinogenesis. atRA supplementation ameliorates colon carcinogenesis in a CD8⁺ T cell-dependent manner.

Highlights

- atRA deficiency in colon cancer is driven by microbiota-induced inflammation
- Treatment with atRA reduces tumorigenesis by enhancing a CD8⁺ T cell response
- The enhanced CD8⁺ T cell response is due to upregulation of MHC I on tumor cells
- CD8⁺ T cell cytotoxicity in human colon cancer correlates with local atRA metabolism



Normalizing Microbiota-Induced Retinoic Acid Deficiency Stimulates Protective CD8⁺ T Cell-Mediated Immunity in Colorectal Cancer

Nupur Bhattacharya,^{1,6,*} Robert Yuan,^{1,6} Tyler R. Prestwood,^{1,2} Hweixian Leong Penny,¹ Michael A. DiMaio,³ Nathan E. Reticker-Flynn,¹ Charles R. Krois,⁴ Justin A. Kenkel,¹ Tho D. Pham,¹ Yaron Carmi,¹ Lorna Tolentino,¹ Okmi Choi,¹ Reyna Hulett,¹ Jinshan Wang,⁴ Daniel A. Winer,⁵ Joseph L. Napoli,⁴ and Edgar G. Engleman^{1,7,*}

¹Department of Pathology, Stanford University School of Medicine (Blood Center), 3373 Hillview Avenue, Palo Alto, CA 94304, USA

²Institute for Stem Cell Biology and Regenerative Medicine, Stanford University School of Medicine, 265 Campus Drive, Stanford, CA 94305, USA

³Department of Pathology, Stanford University, 300 Pasteur Drive, Stanford, CA 94305, USA

⁴Graduate Program in Metabolic Biology, Department of Nutritional Sciences and Toxicology, University of California, Berkeley, Berkeley, CA 94720, USA

⁵Department of Pathology, University Health Network, and Departments of Laboratory Medicine and Pathobiology, and Immunology, University of Toronto, Toronto, ON M5G 2N2, Canada

⁶Co-first author

⁷Lead Contact

*Correspondence: nupur26@stanford.edu (N.B.), edgareng@stanford.edu (E.G.E.)

<http://dx.doi.org/10.1016/j.immuni.2016.08.008>

SUMMARY

Although all-*trans*-retinoic acid (atRA) is a key regulator of intestinal immunity, its role in colorectal cancer (CRC) is unknown. We found that mice with colitis-associated CRC had a marked deficiency in colonic atRA due to alterations in atRA metabolism mediated by microbiota-induced intestinal inflammation. Human ulcerative colitis (UC), UC-associated CRC, and sporadic CRC specimens have similar alterations in atRA metabolic enzymes, consistent with reduced colonic atRA. Inhibition of atRA signaling promoted tumorigenesis, whereas atRA supplementation reduced tumor burden. The benefit of atRA treatment was mediated by cytotoxic CD8⁺ T cells, which were activated due to MHC I upregulation on tumor cells. Consistent with these findings, increased colonic expression of the atRA-catabolizing enzyme, CYP26A1, correlated with reduced frequencies of tumoral cytotoxic CD8⁺ T cells and with worse disease prognosis in human CRC. These results reveal a mechanism by which microbiota drive colon carcinogenesis and highlight atRA metabolism as a therapeutic target for CRC.

INTRODUCTION

Colorectal cancer (CRC) is the second leading cause of cancer mortality in the United States (Haggard and Boushey, 2009), and ulcerative colitis (UC), a chronic inflammatory condition of the colon, has been shown to predispose individuals to CRC (Ullman and Itzkowitz, 2011). Despite advances in therapy, however, 20%–30% of UC patients still undergo colectomy because

they are refractory to current therapy or because they have developed CRC. Unfortunately, surgery is often associated with significant post-operative morbidities (Biondi et al., 2012). Thus, there remains an urgent need for improved therapy and effective chemoprophylaxis in UC and UC-associated cancer.

The vitamin A metabolite all-*trans*-retinoic acid (atRA) is required for several crucial physiological processes (Clagett-Dame and DeLuca, 2002; Mark et al., 2006; Obrochta et al., 2015). In recent years, atRA has been shown to regulate both the innate and adaptive immune systems and, in particular, to play a requisite role in shaping intestinal immunity (Cassani et al., 2012; Hall et al., 2011b). atRA maintains immune homeostasis in the intestinal lamina propria, mainly by potentiating the induction and maintenance of regulatory T cells and reciprocally inhibiting the development of Th17 cells (Benson et al., 2007; Cassani et al., 2012; Coombes et al., 2007; Mucida et al., 2007). Additionally, in certain pathological settings, atRA can also elicit proinflammatory effector T cell responses (Allie et al., 2013; Guo et al., 2012, 2014; Hall et al., 2011a). However, despite the critical influence of atRA on intestinal immunity, its role in CRC has not been previously investigated.

We hypothesized that a local deficiency of atRA might promote the development of CRC, especially in the context of intestinal inflammation. Therefore, we studied atRA metabolism in colitis-associated CRC. Our findings reveal a link between microbiota-induced intestinal inflammation, atRA deficiency, and CRC in mice and humans, as well as a strong anti-tumor effect of atRA mediated through CD8⁺ effector T cells.

RESULTS

Mice with Colitis-Associated Cancer Are Deficient in Colonic atRA due to Altered atRA Metabolism

To investigate the role of atRA metabolism in CRC development, we used a mouse model that recapitulates progression from




Figure 1. AOM-DSS Mice Have a Deficiency in Colonic atRA that Can Be Attributed to Alterations in atRA Metabolism

(A) Mass spectrometry measurements of colonic atRA concentrations in AOM-DSS mice throughout disease progression compared to normal age-matched mice (n = 3–5 mice per group).

(B and C) Immunoblots for ALDH1A1 in distal colonic lysates of mice with chronic colitis (DSS week 7) (B) and of AOM-DSS mice at different time points throughout disease progression (C) compared to normal age-matched mice (n = 5 mice per group). Representative of two independent experiments. Cropped immunoblots were taken from different parts of the same gel and are indicated by demarcation lines.

(D) Immunofluorescence staining of ALDH1A1 (red) in colonic tissue sections throughout disease progression, from steady state to colitis (DSS treated), dysplasia, and carcinoma. Scale bar, 50 μm. n = 5 mice per group.

(legend continued on next page)

colitis to cancer: the AOM-DSS model (Tanaka et al., 2003). In this model, the colonotropic carcinogen azoxymethane (AOM) is combined with the inflammatory agent dextran sodium sulfate (DSS) to induce chronic intestinal inflammation and tumor formation in the distal colons of mice within 9 to 10 weeks, with dysplasia appearing as early as week three. In contrast, mice administered DSS alone develop chronic colitis without tumorigenesis (Wirtz et al., 2007).

In line with our hypothesis that a local deficiency of atRA might promote the development of CRC, colonic atRA levels, as measured by quantitative mass spectrometry, were significantly reduced in mice with colitis-associated cancer (CAC) as early as 4 weeks after AOM-DSS induction, when mice had chronic inflammation with dysplastic changes in the colon. atRA levels further declined to approximately half the normal colonic atRA level by week nine, when carcinomas became apparent (Figures 1A and S1A). To investigate whether this deficiency could result from altered expression of atRA metabolic enzymes, we analyzed the colonic expression of key enzymes that function in the synthesis of atRA—the retinaldehyde dehydrogenases ALDH1A1, ALDH1A2, and ALDH1A3—and the catabolism of atRA—the cytochrome p450 (CYP) family members CYP26A1, CYP26B1, and CYP26C1—during progression from colitis to cancer. Because ALDH1A1 is the most abundantly expressed ALDH1A isoform in the normal mouse colon (data not shown) and CYP26A1 is the most catalytically active of the three CYP26 enzymes (Kedishvili, 2013), we focused our analyses on these enzymes. Colonic ALDH1A1 protein expression declined in mice with chronic colitis and in mice with CAC throughout disease progression, ultimately decreasing by 50%–70% compared to age-matched normal mice (Figures 1B–1D, S1B, and S1C). This reduction was also observed at the transcript level (Figures 1E and 1F). ALDH1A2 protein expression remained unchanged (Figure S1D), whereas ALDH1A3 was consistently decreased during disease progression (Figure S1E).

In contrast to the atRA synthesis enzymes, the colonic transcript levels of the atRA catabolizing enzyme CYP26A1 were increased 4-fold in mice with chronic colitis (Figure 1G), and 3- to 8-fold (depending on the CYP26 isoform) in mice with CAC. Upregulation was observed as early as 1 week after AOM-DSS induction, with levels declining thereafter (Figures 1H, S1F, and S1G).

We next sought to partition out the contribution of various cell types to the colonic atRA deficiency. Although *Aldh1a1* expression was reduced in colonic epithelial cells, T cells, and macrophages or dendritic cells in CAC mice (Figure 1I), *Cyp26a1* expression was increased only in the epithelial compartment (Figure 1J). Given the relative abundance of epithelial cells compared to immune cells (Figure 1K) and the observation that epithelial cells exhibit alterations in both *Aldh1a1* and *Cyp26a1*, the atRA deficiency observed in colitis or CAC can largely be

attributed to the epithelial cells. Taken together, these findings indicate that mice with CAC have a marked decrease in colonic atRA as a result of a decrease in the atRA-synthesizing ALDH1A enzymes and an increase in the atRA-catabolizing CYP26 enzymes.

Abnormal atRA Metabolism Is a Common Feature of Ulcerative Colitis and Colorectal Cancer in Humans

To assess whether atRA deficiency is present in UC-associated CRC, we examined the expression of ALDH1A1 and CYP26A1 in colonic resections from UC and UC-associated CRC patients. In specimens from UC patients, there was a significant decrease in ALDH1A1 and a corresponding increase in CYP26A1 in colitis regions compared to matched uninvolved regions of tissue (Figures 2A and 2B) and in dysplastic tissue compared to matched colitis tissue (Figures 2C and 2D). ALDH1A1 and CYP26A1 were also lower and higher, respectively, in colon adenocarcinomas compared to matched normal regions (Figures 2E and 2F). Using a tissue microarray, we found that the same ALDH1A1 and CYP26A1 differential expression was conserved in sporadic adenocarcinomas compared to normal colons and adenomas (Figures 2G and 2H). Transcript expression of *ALDH1A1* was also decreased and *CYP26A1*, increased, in colon cancer specimens from an independent microarray dataset, GEO: GSE39582 (Figure 2I) (Marisa et al., 2013).

These findings show that patients with sporadic CRC, UC, and UC-associated CRC have similar alterations in atRA metabolism in affected tissues, consistent with the reduced colonic levels of atRA observed in the mouse model of CAC.

Inflammation Triggered by Intestinal Microbiota Induces atRA Enzyme Alteration in Mice with CAC


To investigate whether atRA deficiency can exacerbate CRC tumorigenesis, we treated mice with the pan-retinoic acid receptor (RAR) inverse agonist BMS493 (Germain et al., 2009) 4 weeks prior to and throughout disease progression following AOM-DSS induction. This treatment led to a significant increase in tumor burden (Figure 3A), compelling us to investigate the mechanism responsible for the atRA enzyme deregulation observed in mice with CAC. Several investigators have demonstrated the protumorigenic effects of intestinal microbiota in CRC (Arthur et al., 2012; Couturier-Maillard et al., 2013; Hu et al., 2013), with broad-spectrum antibiotics dramatically reducing tumor incidence in animal models of CRC, including the AOM-DSS model (Grivennikov et al., 2012; Zackular et al., 2013). We therefore hypothesized that bacterial influx into the colon, due to defective barrier permeability, could induce colonic atRA enzyme alteration in the context of colitis and CRC. In accord with our hypothesis, pretreatment with broad-spectrum antibiotics completely prevented both the decrease in *Aldh1a1* and increase in *Cyp26a1* in mice with CAC (Figures 3B, S2A, and S2B).

(E–H) qRT-PCR measurements of *Aldh1a1* (E and F) and *Cyp26a1* (G and H), relative to *Gapdh*, in distal colons of mice treated with DSS (E and G) or AOM-DSS (F and H) compared to normal age-matched mice ($n = 4$ –5 mice per group). Representative of three independent experiments.

(I and J) qRT-PCR measurements of *Aldh1a1* (I) and *Cyp26a1* (J), relative to *Gapdh*, in different cell types sorted from AOM-DSS and normal age-matched mice ($n = 5$ mice per group).

(K) Flow cytometry data showing the frequencies of different cell types in distal colons of AOM-DSS mice ($n = 5$ mice per group). Results are represented as mean \pm SEM; $p < 0.05 = *$; $p < 0.01 = **$; $p < 0.001 = ***$, Mann-Whitney U test.

See also Figure S1.



(legend on next page)

Antibiotic-treated AOM-DSS mice also exhibited significantly increased body weights and dramatically reduced tumor burdens compared to untreated AOM-DSS mice (Figures S2C and S2D).

Having demonstrated that atRA enzymes are altered not only in CAC but also during chronic colitis (Figure 1), we hypothesized that inflammation driven by gut microbiota could contribute to the altered enzyme expression. In line with our hypothesis, the colons of antibiotic-treated AOM-DSS mice not only had healthier colitis scores compared to untreated AOM-DSS mice (Figures 3C and S2E) but also showed a striking reduction in the expression of several proinflammatory cytokines (TNF α , IFN γ , IL-1 β , IL-17A, IL-12, IL-6, and IL-23) (Figure S2F). To test whether these proinflammatory cytokines could be altering atRA enzymes in colitis or CAC, we asked whether culturing epithelial organoids with these cytokines or whether injecting these cytokines into the mucosal wall of the distal colon of antibiotic-treated mice (Figure S2G) could alter the expression of *Aldh1a1* and *Cyp26a1*. In both contexts, the proinflammatory cytokines induced a significant decrease in *Aldh1a1* and increase in *Cyp26a1* (Figures 3D and 3E).

Next, we investigated the cell types responsible for secreting the various proinflammatory cytokines in the colons of mice with CAC. The myeloid compartment secreted a wide range of these cytokines, including TNF α , IL-1 β , IL-6, and IL-12, while the epithelial cells secreted TNF α and T cells secreted IL-17A (Figures 3F and S2H). Antibiotic treatment not only markedly reduced cytokine secretion from all these cell types but also substantially reduced immune cell infiltration into the colon (Figures 3F, 3G, and S2H). These results show that expression of the atRA metabolic enzymes is altered by inflammatory cytokines released from colonic epithelial cells and infiltrating immune cells in response to the influx of gut microbiota across a defective mucosal barrier.

atRA Supplementation Decreases Tumor Burden in Mice with CAC

Given the altered atRA metabolism in AOM-DSS mice, as well as in human CRC, we assessed whether treatment with atRA could decrease tumor burden. We used two distinct methods for this purpose: (1) intraperitoneal (i.p.) injection of atRA and (2) oral administration of Liarozole, an inhibitor of the CYP26 enzymes that catabolize atRA (Van Wauwe et al., 1990). Liarozole's ability to increase available endogenous atRA was demonstrated in vitro using RARE-luciferase assays (Figures S3A and S3B) on two colon cancer cell lines, MC38 and CT26, and in vivo via mass spectrometry of colons of AOM-DSS mice orally gavaged with Liarozole (Figure S3C). atRA was administered to AOM-DSS mice either starting early, when acute colitis was established, or

later, after dysplastic changes had occurred in the colon. In both cases, tumor burden in atRA-treated mice was reduced by more than half compared to vehicle-treated mice (Figures 4A and 4B). Consistent with atRA treatment, Liarozole treatment also resulted in a 50% reduction in tumor burden (Figure 4C). These data demonstrate a robust therapeutic benefit conferred by atRA supplementation.

atRA Treatment Elicits a Pronounced CD8⁺ T Cell Response in Mice with CAC

Next, we sought to identify the mechanism responsible for the anti-tumor effect of atRA. Since CD8⁺ T cells play a crucial role in anti-tumor immunity, we examined the CD8⁺ T cell response in AOM-DSS mice treated with atRA. A significant increase in the percentage of CD8⁺ T cells expressing the early activation marker CD69 was seen in tumors and mesenteric lymph nodes (MLNs) in atRA-treated mice (Figures 5A, 5B, and S4A). The atRA-mediated increase in activated CD8⁺ T cells was more pronounced in the lamina propria (LP) than in the intra-epithelial lymphocyte (IEL) layer of the tumor (Figure 5C). In addition, the percentage of proliferating intratumoral CD8⁺ T cells nearly doubled in atRA-treated mice (Figure 5D). Thus, a robust CD8⁺ T cell response is induced by atRA treatment of mice with CAC.

The Anti-tumor Effect of atRA Is CD8⁺ T Cell Dependent

To determine the functional relevance of the CD8⁺ T cell response elicited by atRA treatment, we evaluated the therapeutic benefit of atRA in AOM-DSS mice depleted of CD8⁺ T cells (Figure S4B), as well as in CD8-deficient (*Cd8a*^{-/-}) mice. atRA had no effect on tumor burden in either CD8⁺ T cell-depleted mice or *Cd8a*^{-/-} mice (Figures 5E and 5H), demonstrating that the therapeutic effect of atRA is dependent on CD8⁺ T cells. Consistent with this idea, we observed a 4-fold increase in tumor cell death in atRA-treated AOM-DSS mice compared to vehicle-treated mice (Figures 5F and 5G), with CD8⁺ T cells in close proximity to TUNEL⁺ tumor cells in the atRA-treated condition (Figure S4C). This increase in tumor cell death was completely abrogated in CD8⁺ T cell-depleted mice (Figures 5F and 5G). Importantly, the therapeutic benefit of atRA treatment in *Cd8a*^{-/-} mice was restored upon adoptive transfer of CD8⁺ T cells (Figures 5H and S4D). Although mature CD4⁺ T cells can be reprogrammed in an atRA-dependent manner into MHCII-restricted CD8 $\alpha\alpha$ ⁺ IELs with cytotoxic functions (Mucida et al., 2013; Reis et al., 2014), in the context of CAC, atRA treatment did not increase the frequency of this cell type (Figure S4E). These results therefore demonstrate that the therapeutic benefit conferred by atRA is dependent on CD8⁺ T cells.


A detailed analysis of CD4⁺ T cells did not reveal any significant changes in the number (Figure S5A) or activation status of

Figure 2. Human UC, UC-Associated Dysplasia, and Sporadic Colon Adenocarcinoma Exhibit Abnormal Expression of Colonic atRA-Metabolizing Enzymes

(A, C, E, and G) Representative immunofluorescence images of ALDH1A1 and CYP26A1 (red) on matched human UC and uninvolved colonic resections (n = 6 samples) (A), matched human UC and dysplastic colonic resections (n = 6 samples) (C), matched adenocarcinoma and normal colonic resections (n = 5 samples) (scale bar, 50 μ m) (E), and on a sporadic adenoma and carcinoma tissue microarray (G) (scale bar, 1 mm).

(B, D, F, and H) Average staining intensity values (in a.u.) of ALDH1A1 and CYP26A1 in the colonic epithelium, acquired from a minimum of three image fields, of all samples in (A, C, E, and G).

(I) Transcript expression of *ALDH1A1* and *CYP26A1* from the gene expression microarray dataset GEO: GSE39582. Paired t test (B, D, and F) and one-way ANOVA test (H), p < 0.05 = *; p < 0.01 = **; p < 0.001 = ***.



(legend on next page)

these cells in the tumors and MLNs of atRA-treated mice (Figures S5B and S5C). Moreover, CD4⁺ T cells isolated from the tumor or surrounding tissue from atRA-treated mice did not show any differences in TNF α , IFN γ , or IL-17A secretion compared to vehicle-treated mice (Figures S5D and S5E). Interestingly, there was a significant increase in the frequency of CD4⁺FOXP3⁺ regulatory T (Treg) cells in tumors of atRA-treated mice (Figure S5F); however, this increase in tumoral Treg cells was also observed in atRA-treated *Cd8a*^{-/-} mice (Figure S5G), which are refractory to the therapeutic benefit of atRA, indicating that the anti-tumor effect of atRA is not mediated through Treg cells alone.

In a study in mice with DSS-induced colitis, atRA supplementation suppressed colitis by inducing $\gamma\delta$ ⁺ T cells to secrete IL-22 (Mielke et al., 2013). However, in our studies, atRA treatment of T cell receptor delta chain (*Tcrd*^{-/-}) mice still significantly decreased tumor burden, indicating that $\gamma\delta$ ⁺ T cells are not required for the anti-tumor effect of atRA (Figure S5H). Nonetheless, we observed that the colonic tissue surrounding the tumoral regions in atRA-treated mice had reduced inflammation scores, which could contribute to the atRA-mediated reduction in tumor burden in parallel with the CD8⁺ T cell mechanism (Figures S5I and S5J).

atRA Upregulates MHC I Expression on Tumor Epithelial Cells to Promote T Cell Cytotoxicity

Given that CD8⁺ T cells are the dominant mediator of the anti-tumor effect of atRA, we sought to determine the mechanistic basis of the CD8⁺ T cell-dependent effect of atRA treatment on CAC. Since CD8⁺ T cell-mediated cytotoxicity is MHC class I restricted, we analyzed the expression of MHC I on the tumor epithelial cells before and after atRA treatment and found that mice treated with atRA exhibited a significant increase in MHC I (Figures 6A and S6A). Interestingly, this atRA-mediated increase in MHC I was abrogated when the RAR α was deleted from intestinal epithelial cells using *Vil1-cre-Rara*^{fl/fl} mice (Figure 6B). In line with the direct mode of action of atRA on tumor epithelial cells, exogenous addition of atRA to the human colon cancer cell line Caco-2 in vitro also increased MHC I expression (Figure 6C).

Next, we considered the possibility that increased MHC I expression on tumor epithelial cells could render them more susceptible to CD8⁺ cytotoxic T cell killing. Indeed, *Vil1-cre-Rara*^{fl/fl} mice that lack the atRA-mediated upregulation of MHC I were resistant to the anti-tumor effects of atRA (Figure 6D). Consistent with this, intratumoral CD8⁺ T cells in atRA-treated AOM-DSS

wild-type (WT) mice expressed higher levels of granzyme B, indicating increased cytotoxic activity (Figures 6E and S6B), which was not seen in *Vil1-cre-Rara*^{fl/fl} mice (Figure 6F). Consequently, untreated *Vil1-cre-Rara*^{fl/fl} mice induced with AOM-DSS had a trend toward a higher tumor burden than control mice (Figure S6C). To determine if atRA acted on epithelial cells—and not myeloid cells or CD8⁺ T cells—to mediate its anti-tumor effects, we tested whether mice lacking RAR α expression in macrophages (*Lyz2-cre-Rara*^{fl/fl} mice), dendritic cells (*Ilgax-cre-Rara*^{fl/fl} mice), and CD8⁺ T cells (through adoptive transfer of CD8⁺ T cells from *Cd4-cre-Rara*^{fl/fl} mice into *Cd8a*^{-/-} mice, as CD8⁺ T cells in these mice lack the receptor due to cre expression in the double-positive thymocytes) were responsive to the anti-tumor effects of atRA. Importantly, all of the aforementioned mice exhibited lower tumor burdens upon atRA treatment, confirming that the anti-tumor effect of atRA was not dependent on direct effects of atRA on these cell types (Figures S6D–S6F). These results demonstrate that atRA acts directly on tumor epithelial cells to upregulate MHC I, thereby rendering them sensitive to CD8⁺ T cell-mediated killing.

To assess our findings in another model of CRC, we investigated whether the MC38 subcutaneous tumor model was similarly responsive to atRA through a CD8⁺ T cell-mediated mechanism. In line with our observations from the AOM-DSS model, we found that, while WT mice subcutaneously implanted with MC38 were responsive to the anti-tumor effect of atRA, *Cd8a*^{-/-} mice were not (Figures S7A and S7B). Moreover, atRA-treated MC38 tumor mice had a significantly higher expression of MHC I on their tumor cells compared to vehicle-treated mice (Figure S7C).

CYP26A1 in Colon Carcinoma Correlates with Reduced Cytotoxic CD8⁺ T Cell Frequency and Worse Disease Prognosis

The findings from our mouse studies suggest that intestinal atRA deficiency dampens the CD8⁺ T cell-mediated anti-tumor responses, which are known to correlate with overall survival in CRC (Deschoolmeester et al., 2010; Galon et al., 2006; Koelzer et al., 2014). We therefore correlated ALDH1A1 and CYP26A1 expression in each core biopsy from the sporadic colon cancer tissue microarray with tumoral CD8⁺ T cell density and granzyme B expression to determine if atRA metabolism could predict cytotoxic T cell potential. Indeed, there was a significant negative correlation between CYP26A1 expression and both tumoral

Figure 3. Inflammation Triggered by Intestinal Microbiota Induces atRA Enzyme Alteration in AOM-DSS Mice

- (A) Tumor analyses in AOM-DSS mice treated with the pan-retinoic acid receptor inhibitor (BMS493) compared to vehicle-treated mice. Also shown are representative images of the colons. Data are pooled from 2 independent experiments (n = 7–8 mice).
- (B) qRT-PCR measurements of *Aldh1a1* and *Cyp26a1*, normalized to *Gapdh* in the distal colons of normal age-matched mice and untreated or antibiotic-treated AOM-DSS mice (n = 5–6 mice per group). Representative of two independent experiments.
- (C) Colitis score grading of antibiotic-treated or untreated AOM-DSS mice (n = 5 mice per group).
- (D) qRT-PCR measurements of *Aldh1a1* and *Cyp26a1* from intestinal organoids cultured in vitro with a proinflammatory cytokine mixture compared to vehicle. Data are pooled from three independent experiments.
- (E) qRT-PCR measurements of *Aldh1a1* and *Cyp26a1* from distal colons of antibiotic-treated mice injected intramucosally with a proinflammatory cytokine mixture or vehicle (n = 8–10 mice per group).
- (F) Flow cytometry data showing the percentage of specific cell populations from the distal colons of untreated or antibiotic-treated AOM-DSS mice expressing different cytokines induced via stimulation with fecal extract (n = 5 mice per group).
- (G) Quantitation of the frequency of different immune cells in the distal colons of normal age-matched mice and untreated or antibiotic-treated AOM-DSS mice (n = 5 mice per group). Results are represented as mean \pm SEM; p < 0.05 = *; p < 0.01 = **; p < 0.001 = ***, Mann-Whitney U test. See also Figure S2.





Figure 4. atRA Supplementation Decreases Tumor Burden in AOM-DSS Mice

(A and B) Graphs represent tumor analyses in AOM-DSS mice treated with 200 μ g atRA twice a week starting from day 10 (A) or every other day from week 4 (B) after disease induction compared to vehicle-treated mice. Representative image of mouse colons from (B). Data represent pooled data from 2 independent experiments with $n = 10$ mice per group in (A) and $n = 9-13$ mice in (B).

(C) Tumor analyses in AOM-DSS mice fed with a diet containing Liarozole compared to a control base diet from day 10 after disease induction. Representative of 3 independent experiments ($n = 10$ mice per group). Results are represented as mean \pm SEM; $p < 0.05 = *$; $p < 0.01 = **$; $p < 0.001 = ***$, Mann-Whitney U test. See also Figure S3.



(legend on next page)

CD8⁺ T cell density (Figures 7A–7C) and percentage of CD8⁺ T cells expressing granzyme B in CRC specimens (Figures 7A, 7B, and 7D). We found no significant correlation with ALDH1A1 (data not shown). We also tested whether *ALDH1A1* and *CYP26A1* mRNA expression correlates with disease prognosis using the GEO: GSE39582 dataset (Marisa et al., 2013). This dataset has a large number of CRC samples obtained from a multi-center cohort (n = 566) containing information on overall survival and disease-free survival. Although *ALDH1A1* expression did not correlate with disease prognosis, *CYP26A1* expression correlated inversely with both overall survival and disease-free survival (Figures 7E and 7F).

DISCUSSION

atRA has shown promise in the treatment of several malignancies. In acute promyelocytic leukemia, atRA is known to act by inducing post-maturation apoptosis of the leukemic cells (Jiménez-Lara et al., 2004). Despite its established regulatory role in intestinal immunity, the influence of atRA on CRC development has not been previously examined. Our findings reveal that mice with CAC have significantly reduced levels of atRA in their colons due to a marked decrease in atRA-synthesizing ALDH1A enzymes and an increase in atRA-catabolizing CYP26 enzymes, imputing an important role to atRA in CRC.

Direct evidence of the adverse effect of an atRA deficit was demonstrated in mice with CAC. In these mice, further inhibition of atRA signaling increased tumor burden, while, conversely, atRA supplementation reduced tumor burden. Interestingly, depletion of the intestinal microbiota in AOM-DSS mice prevented the alteration of the atRA enzymes. The absence of intestinal bacteria in germ-free mice and antibiotic-treated mice has been demonstrated to dramatically reduce tumor formation in several models of CRC (Grivennikov et al., 2012; Zackular et al., 2013). Increased inflammation and production of genotoxic metabolites are among the mechanisms by which microbiota potentiate colon carcinogenesis (Belcheva et al., 2014; Grivennikov, 2013). However, our study demonstrates yet another mechanism by which intestinal microbiota may promote CRC: by altering atRA metabolism.

An important finding of our study is that CRC patients have similar alterations in atRA metabolism in their tumors irrespective of whether they had a history of colitis. High inter-patient variability in our tissue microarray analysis of ALDH1A1 and CYP26A1 in sporadic colon carcinomas and adenomas may

reflect differences in the extent of barrier permeability or associated inflammation. Similarly, adenocarcinomas may exhibit lower expression of epithelial tight junction proteins and greater barrier permeability compared to adenomas, possibly explaining the more pronounced enzyme changes in carcinomas (Nakayama et al., 2008; Soler et al., 1999). Nevertheless, when compared with normal colonic mucosa, colitis, dysplasia, and colon carcinoma are all characterized by a decrease in ALDH1A1 and a concomitant increase in the CYP26A1, indicative of reduced colonic levels of atRA. These findings are in accord with recent investigations of sporadic CRC (Kropotova et al., 2014), including a study demonstrating that CYP26B1 expression directly correlates with disease prognosis (Brown et al., 2014). In the present study, we found a significant correlation between *CYP26A1* transcript expression and both overall survival and disease-free survival in a multi-center cohort of CRC patients (Marisa et al., 2013). Our findings that abnormal atRA metabolism correlates with clinical prognosis and is present across most pre-neoplastic and neoplastic conditions of the colon, combined with the observed exacerbation of disease in mice with CAC upon inhibition of atRA signaling and the therapeutic benefit of atRA supplementation, suggest that atRA deficiency is an important factor in the pathogenesis of CRC.

A number of possible mechanisms might explain the beneficial anti-tumor effect of atRA. In a mouse model of Crohn's disease, atRA supplementation attenuated disease progression by promoting Treg-cell-mediated intestinal tolerance (Collins et al., 2011). In another study, atRA supplementation ameliorated DSS-induced colitis but through a mechanism that was $\gamma\delta^+$ -T cell dependent. (Mielke et al., 2013). However, in our studies, the anti-tumor effect of atRA was found to be independent of $\gamma\delta^+$ T cells. Since in vitro studies have shown that atRA can induce apoptosis of colon cancer cell lines, albeit at high non-physiological concentrations (Bengtsson et al., 2013; Lee et al., 2000), we considered the possibility that the therapeutic benefit of atRA was due to a direct apoptotic effect on the tumor cells. In our studies, atRA treatment did lead to tumor cell apoptosis in vivo, but, surprisingly, the therapeutic effect of atRA was mediated by CD8⁺ T cells rather than through a direct anti-tumor effect.

Previous studies have shown that atRA can elicit CD4⁺ or CD8⁺ effector T cell responses in the context of infection (Guo et al., 2014; Hall et al., 2011b). Here, in CAC, we found that atRA treatment led to an increase in tumoral CD8⁺ T cell cytotoxic activity. Furthermore, using *Vil1-cre-Rara^{fl/fl}* mice, we found

Figure 5. atRA Reduces Tumor Burden in AOM-DSS Mice through a CD8⁺ T Cell-Dependent Mechanism

(A–D) Flow cytometry data of percentages of CD69⁺ CD8⁺ T cells (A–C) and Ki67⁺ CD8⁺ T cells (D) in tumors (A and D), surrounding distal colonic tissue (A and D), and IEL and LP layers of the tumors (C) and MLNs (B) of vehicle- or atRA-treated AOM-DSS mice. Data were pooled from 3 independent experiments with n = 14 mice per group (A) and n = 8 mice per group (B) and from 2 independent experiments with n = 6–10 mice per group (C) and n = 6 mice per group (D).

(E) Tumor analyses of vehicle- or atRA-treated AOM-DSS mice injected from week 4 to week 9 with isotype control or anti-CD8 α depleting antibody. Data were pooled from 2 independent experiments with n = 6 mice in the isotype-treated group and n = 8 mice in the CD8-depleted group.

(F) Immunofluorescence images showing EpCAM (green), DAPI (blue), and TUNEL (red) in colonic tumor sections from atRA- or vehicle-treated isotype-injected and CD8-depleted AOM-DSS mice. Representative image from n = 6 per group in the isotype-treated group and n = 4 mice in the CD8-depleted group. Scale bar, 50 μ m.

(G) Relative number of TUNEL⁺ cells quantified by normalizing to tumor crypt area in (F).

(H) Tumor analyses in *Cd8a^{-/-}* mice induced with AOM-DSS, and *Cd8a^{-/-}* mice adoptively transferred with CD8⁺ T cells, treated from week 3 to week 9 with atRA (n = 6–9 mice per group). Results are represented as mean \pm SEM; p < 0.05 = *; p < 0.01 = **; p < 0.001 = ***, Mann-Whitney U test.

See also Figures S4 and S5.




Figure 6. atRA Upregulates MHC I Expression on Tumor Epithelial Cells, Leading to Increased Cytotoxic T Lymphocytes

(A and B) Flow cytometry plots of geometric mean intensity (MFI) of MHC I expression on tumor and surrounding epithelial cells (EpCAM⁺) from atRA- or vehicle-treated normal age-matched (A) or *Vil1-cre-Rara^{fl/fl}* (B) AOM-DSS mice. Also shown are representative histograms of MHC I from the tumor epithelial cells. Data were pooled from 2 independent experiments with $n = 8-9$ mice per group.

(C) MHC I MFI on Caco-2 cells 72 hr after in vitro treatment with 1 μ M atRA compared to vehicle (DMSO). Shown are three independent experiments and a representative histogram.

(D) Tumor analyses of vehicle- or atRA-treated *Vil1-cre-Rara^{fl/fl}* mice induced with AOM-DSS. Data were pooled from 2 independent experiments, with $n = 8-9$ mice per group.

(E and F) Flow cytometry data of percentage of granzyme B⁺ CD8⁺ T cells in the tumors and surrounding distal colonic tissue from atRA- versus vehicle-treated normal age-matched (E) or *Vil1-cre-Rara^{fl/fl}* (F) AOM-DSS mice. Data were pooled from 2 independent experiments with $n = 6$ mice per group. Results are represented as mean \pm SEM $p < 0.05 = *$; $p < 0.01 = **$; $p < 0.001 = ***$, Mann-Whitney U test.

See also [Figures S6 and S7](#).

that atRA treatment directly acts on tumor epithelial cells to upregulate MHC I expression, thereby rendering them more susceptible to killing by CD8⁺ cytotoxic T lymphocytes. Consistent with this mechanism, previous in vitro studies have demonstrated that MHC I is a direct transcriptional target of atRA (Balmer and Blomhoff, 2002; Jansa and Forejt, 1996; Nagata et al., 1992). Untreated *Vil1-cre-Rara^{fl/fl}* mice induced with AOM-DSS did not develop more tumors than control mice, but since mice with CAC already have dramatically reduced levels

of atRA, the absence of RAR α on epithelial cells likely would have had little impact on tumor burden.

Our findings from the mouse model of CAC suggest that atRA deficiency in human CRC could impair effector cytotoxic T cell function, resulting in accelerated growth of tumors. Consistent with this hypothesis, we found a strong negative correlation between abnormal atRA metabolism and both tumoral CD8⁺ T cell frequency and granzyme B expression in human CRC. Several studies have shown that infiltration of T cells, especially CD8⁺




Figure 7. CYP26A1 in Colon Carcinoma Correlates with Reduced CD8⁺ T Cell Cytotoxicity, as well as with Worse Disease Prognosis

(A and B) Representative images of core biopsies of adenocarcinomas showing high and low CYP26A1 staining (red) and corresponding serial sections showing CD8⁺ T cell staining (green) and granzyme B staining (red). Scale bar, 50 μ m.

(C and D) Correlation between CYP26A1 staining intensity (in a.u.) and CD8⁺ T cell density normalized to tumor area (C) and percentage of CD8⁺ T cells expressing granzyme B (D) in sporadic adenocarcinoma specimens, using Pearson's correlation analyses.

(E and F) Kaplan-Meier curves showing the correlation of CYP26A1 transcript expression, partitioned around the median, with overall survival (E) and disease-free survival (F) in colon cancer patients. $p < 0.05 = *$; $p < 0.01 = **$; $p < 0.001 = ***$.

colon could provide a therapeutic benefit in human CRC by promoting MHC1 expression by tumor cells and cytotoxic T cell function.

EXPERIMENTAL PROCEDURES

Patient Specimens

Patient specimens were obtained from the Stanford Tissue Bank and from the Toronto General Hospital under protocols approved by the Institutional Review Board (IRB, Stanford) and the University Health Network Research Ethics Board (Toronto General Hospital).

Mice

C57BL/6 mice, *Cd8a*^{-/-} mice, *Vil1*-cre mice, *Lyz2*-cre mice, *Cd4*-cre mice, and *Tcrd*^{-/-} mice were from The Jackson Laboratory. *Rara*^{fl/fl} mice were a kind gift from Dr. Yasmine Belkaid (National Institute of Allergy and Infectious Diseases).

DSS and AOM-DSS Mouse Models

The DSS and AOM-DSS mouse models of chronic colitis and CAC, respectively, were established using the protocol described by [Wirtz et al. \(2007\)](#) with slight modifications. In brief, for the development of chronic colitis, mice were given 3% DSS salt in drinking water for 7 days, followed by drinking water for 14 days. This cycle was repeated twice. For CAC, mice were given an initial i.p. injection of AOM at the beginning of the chronic colitis protocol.

Antibiotic Treatment of AOM-DSS Mice

Antibiotics were supplied in water before induction with AOM-DSS. The antibiotics used were ampicillin (1g/L), metronidazole (1g/L), vancomycin (0.5 g/L), and neomycin (1g/L) (Sigma-Aldrich).

Bacterial Culture on Agar Plates

Fecal extract from antibiotic-treated mice was cultured on Luria broth (LB) Agar and Blood Agar (TSA with 5% sheep blood) plates.

T cells, as well as MHC1 expression by tumor cells, correlates with improved CRC prognosis ([Deschoolmeester et al., 2010](#); [Galon et al., 2006](#); [Koelzer et al., 2014](#); [Simpson et al., 2010](#)). Our work not only provides a potential explanation for these phenomena but also suggests that restoration of atRA levels in the

atRA Treatment of Caco-2 Cell Line

Caco-2 cells were cultured in complete Eagle's Minimum Essential Medium (EMEM) medium and treated with 1 μ M atRA or DMSO. They were harvested for analysis 72 hr later.

Luciferase Assay

CT26 or MC38 cells were transfected with a firefly luciferase reporter plasmid driven by a pGL3-RARE-responsive promoter (Addgene) and a renilla luciferase reporter plasmid. atRA or vehicle was added along with increasing amounts of Liarozole, and the activity of both reporters was recorded.

Intramucosal Colonic Injections

Antibiotic-treated C57BL/6 mice were injected intramucosally in the distal colon with a cytokine mixture containing 100 ng each of TNF α , IL-1 β , IL-12p70, IL-23, IL-6, IFN γ , and IL-17A (Peprotech) or 1% BSA in PBS (vehicle), and the distal colonic tissues were collected for qRT-PCR analyses.

Intestinal Organoid Culture and Cytokine Treatment

Organoids derived from tumors of AOM-DSS mice were cultured with a cytokine mixture containing 50 ng/mL each of TNF α , IL-1 β , IL-12p70, IL-23, IL-6, and IFN γ or 1% BSA in PBS (vehicle) for 48 hr and then harvested for qRT-PCR analyses.

Quantitation of All-trans-Retinoic Acid Expression in Mouse Colonic Tissue

atRA was extracted from mouse colons as described (Kane et al., 2008). Quantification of retinoids was done using liquid chromatography-tandem mass spectrometry (LC-MS/MS) with an alternate LC separation.

Drug Treatment

200 μ g BMS493 (Tocris Bioscience) or DMSO alone (vehicle) was i.p. injected into female C57BL/6 mice 4 weeks prior to induction with AOM-DSS and continued until the end of the AOM-DSS treatment period. atRA (200 μ g) or DMSO alone (vehicle) was i.p. injected into AOM-DSS mice either twice a week or every other day. Liarozole (80 ppm) (Tocris Bioscience) was incorporated into a base diet containing 4 IU/g of vitamin A (Research Diets). Mice were orally gavaged with 400 μ g Liarozole or polyethylene glycol-200 (vehicle) after induction with AOM-DSS daily until the end of week 4.

CD8⁺ T Cell Depletion

500 μ g rat anti-mouse CD8 α antibody (clone YTS169.4, BioXCell) was injected i.p. into 10-week-old C57BL/6 female mice once a week, from week 4 to week 9 after disease induction. The control group received the same amount of isotype control antibody (clone LTF2, BioXCell).

Adoptive Transfer of CD8⁺ T Cells into *Cd8a*^{-/-} Mice

CD8⁺ T cells were isolated from the spleen, MLNs, and other lymph nodes from DSS-treated C57BL/6 mice and adoptively transferred into *Cd8a*^{-/-} mice (8×10^6 cells per mouse) 2 weeks after induction with AOM-DSS (Figure S4D). atRA or vehicle was administered every other day to recipient mice from the day of adoptive transfer to the end of week 9.

Immunoblotting

Protein lysates were prepared from the distal half of the colons using RIPA lysis buffer. Primary and secondary HRP-labeled antibodies were used at 1:500 and 1:2,000 dilutions, respectively, in 5% milk in PBS-Tween.

qRT-PCR

Transcript expression was determined using the Fast SYBR Green PCR Master Mix (Applied Biosystems) and the 7900HT real-time PCR instrument (Applied Biosystems). For quantification of 16S rDNA, mouse fecal DNA was extracted using the REExtract-N-Amp Tissue PCR Kit (Sigma Aldrich) and purified of inhibitors (Zymo Research).

Histology and Colitis Grading

Disease diagnosis and colitis grading of H&E-stained slides were determined by an experienced gastrointestinal pathologist at Stanford University who was

blinded with respect to treatment. For colitis grading, the scheme from Geboes et al. (2000) was used.

Immunofluorescence Staining

OCT-embedded colonic sections were post-fixed in 2% paraformaldehyde and blocked with 10% goat serum. They were subsequently stained with primary and secondary antibodies.

Flow Cytometry

Isolated colonic and MLN cells were resuspended in a fluorescence-activated cell sorting (FACS) buffer containing 1% BSA in PBS and stained with antibodies. Flow cytometric data acquisition was performed on a LSRII flow cytometer (BD Biosciences).

Statistics

Experimental data were analyzed with the Mann-Whitney U test using Prism (GraphPad Software), unless otherwise stated. Tissue microarray data were analyzed using the one-way ANOVA test, and correlation analysis was performed using the Pearson's correlation test. Kaplan Meier curves for disease-free survival and overall survival were generated and analyzed with Prism (GraphPad Software) using the log-rank (Mantel Cox) test. Results are represented as mean \pm SEM; $p < 0.05 = *$; $p < 0.01 = **$; $p < 0.001 = ***$.

Animal Study Approval

All animal protocols used in this study were approved by the Stanford University Institutional Animal Care and Use Committee.

SUPPLEMENTAL INFORMATION

Supplemental Information includes Supplemental Experimental Procedures and seven figures and can be found with this article online at <http://dx.doi.org/10.1016/j.immuni.2016.08.008>.

AUTHOR CONTRIBUTIONS

N.B. and R.Y. conceived the study, performed all experiments, and wrote the manuscript. T.R.P., H.L.P., N.E.R.-F., J.A.K., and Y.C. helped in designing the research and provided technical assistance. M.A.D. and T.D.P. provided pathology assistance. R.H. provided experimental help. C.R.K., J.W., and J.L.N. provided experimental assistance for the atRA mass spectrometry measurements. L.T. and O.C. provided technical assistance with flow cytometry. D.A.W. provided patient specimens for histology. E.G.E. supervised the study and wrote the manuscript. All authors read and approved the final manuscript.

ACKNOWLEDGMENTS

We thank Professor Robert West for providing us with colon cancer specimens. We thank Raghav Kumar for technical assistance, Nancy Wu for flow cytometry assistance, and Philip Engleman and Shawn Winer for help with the pathological diagnoses of histology slides. We also wish to thank Matt Spitzer and Ian Linde for critical review of the manuscript. This work was supported by NIH grants 5 U01 CA141468 and 1 R01 CA163441. T.R.P. is supported by NIH NRSA F30CA196145. N.E.R.-F. is supported by NIH NRSA F32CA189408.

Received: July 21, 2015

Revised: April 7, 2016

Accepted: June 6, 2016

Published: August 30, 2016

REFERENCES

- Allie, S.R., Zhang, W., Tsai, C.Y., Noelle, R.J., and Usherwood, E.J. (2013). Critical role for all-trans retinoic acid for optimal effector and effector memory CD8 T cell differentiation. *J. Immunol.* 190, 2178–2187.
- Arthur, J.C., Perez-Chanona, E., Mühlbauer, M., Tomkovich, S., Uronis, J.M., Fan, T.J., Campbell, B.J., Abujamel, T., Dogan, B., Rogers, A.B., et al. (2012).

- Intestinal inflammation targets cancer-inducing activity of the microbiota. *Science* 338, 120–123.
- Balmer, J.E., and Blomhoff, R. (2002). Gene expression regulation by retinoic acid. *J. Lipid Res.* 43, 1773–1808.
- Belcheva, A., Irrazabal, T., Robertson, S.J., Streutker, C., Maughan, H., Rubino, S., Moriyama, E.H., Copeland, J.K., Kumar, S., Green, B., et al. (2014). Gut microbial metabolism drives transformation of MSH2-deficient colon epithelial cells. *Cell* 158, 288–299.
- Bengtsson, A.M., Jönsson, G., Magnusson, C., Salim, T., Axelsson, C., and Sjölander, A. (2013). The cysteinyl leukotriene 2 receptor contributes to all-trans retinoic acid-induced differentiation of colon cancer cells. *BMC Cancer* 13, 336.
- Benson, M.J., Pino-Lagos, K., Roseblatt, M., and Noelle, R.J. (2007). All-trans retinoic acid mediates enhanced T reg cell growth, differentiation, and gut homing in the face of high levels of co-stimulation. *J. Exp. Med.* 204, 1765–1774.
- Biondi, A., Zoccali, M., Costa, S., Troci, A., Contessini-Avesani, E., and Fichera, A. (2012). Surgical treatment of ulcerative colitis in the biologic therapy era. *World J. Gastroenterol.* 18, 1861–1870.
- Brown, G.T., Cash, B.G., Bliohoghe, D., Johansson, P., Alnabulsi, A., and Murray, G.I. (2014). The expression and prognostic significance of retinoic acid metabolising enzymes in colorectal cancer. *PLoS ONE* 9, e90776.
- Cassani, B., Villablanca, E.J., De Calisto, J., Wang, S., and Mora, J.R. (2012). Vitamin A and immune regulation: role of retinoic acid in gut-associated dendritic cell education, immune protection and tolerance. *Mol. Aspects Med.* 33, 63–76.
- Clagett-Dame, M., and DeLuca, H.F. (2002). The role of vitamin A in mammalian reproduction and embryonic development. *Annu. Rev. Nutr.* 22, 347–381.
- Collins, C.B., Aherne, C.M., Kominsky, D., McNamee, E.N., Lebsack, M.D., Eitzschig, H., Jedlicka, P., and Rivera-Nieves, J. (2011). Retinoic acid attenuates ileitis by restoring the balance between T-helper 17 and T regulatory cells. *Gastroenterology* 141, 1821–1831.
- Coombes, J.L., Siddiqui, K.R., Arancibia-Cárcamo, C.V., Hall, J., Sun, C.M., Belkaid, Y., and Powrie, F. (2007). A functionally specialized population of mucosal CD103+ DCs induces Foxp3+ regulatory T cells via a TGF- β and retinoic acid-dependent mechanism. *J. Exp. Med.* 204, 1757–1764.
- Couturier-Maillard, A., Secher, T., Rehman, A., Normand, S., De Arcangelis, A., Haesler, R., Huot, L., Grandjean, T., Bressenot, A., Delanoye-Crespin, A., et al. (2013). NOD2-mediated dysbiosis predisposes mice to transmissible colitis and colorectal cancer. *J. Clin. Invest.* 123, 700–711.
- Deschoolmeester, V., Baay, M., Van Marck, E., Weyler, J., Vermeulen, P., Lardon, F., and Vermorken, J.B. (2010). Tumor infiltrating lymphocytes: an intriguing player in the survival of colorectal cancer patients. *BMC Immunol.* 11, 19.
- Galon, J., Costes, A., Sanchez-Cabo, F., Kirilovsky, A., Mlecnik, B., Lagorce-Pagès, C., Tosolini, M., Camus, M., Berger, A., Wind, P., et al. (2006). Type, density, and location of immune cells within human colorectal tumors predict clinical outcome. *Science* 313, 1960–1964.
- Geboes, K., Riddell, R., Ost, A., Jensfelt, B., Persson, T., and Löfberg, R. (2000). A reproducible grading scale for histological assessment of inflammation in ulcerative colitis. *Gut* 47, 404–409.
- Germain, P., Gaudon, C., Pogenberg, V., Sanglier, S., Van Dorsselaer, A., Royer, C.A., Lazar, M.A., Bourguet, W., and Gronemeyer, H. (2009). Differential action on coregulator interaction defines inverse retinoid agonists and neutral antagonists. *Chem. Biol.* 16, 479–489.
- Grivennikov, S.I. (2013). Inflammation and colorectal cancer: colitis-associated neoplasia. *Semin. Immunopathol.* 35, 229–244.
- Grivennikov, S.I., Wang, K., Mucida, D., Stewart, C.A., Schnabl, B., Jauch, D., Taniguchi, K., Yu, G.Y., Osterreicher, C.H., Hung, K.E., et al. (2012). Adenoma-linked barrier defects and microbial products drive IL-23/IL-17-mediated tumour growth. *Nature* 491, 254–258.
- Guo, Y., Pino-Lagos, K., Ahonen, C.A., Bennett, K.A., Wang, J., Napoli, J.L., Blomhoff, R., Sockanathan, S., Chandraratna, R.A., Dmitrovsky, E., et al. (2012). A retinoic acid-rich tumor microenvironment provides clonal survival cues for tumor-specific CD8(+) T cells. *Cancer Res.* 72, 5230–5239.
- Guo, Y., Lee, Y.C., Brown, C., Zhang, W., Usherwood, E., and Noelle, R.J. (2014). Dissecting the role of retinoic acid receptor isoforms in the CD8 response to infection. *J. Immunol.* 192, 3336–3344.
- Haggar, F.A., and Boushey, R.P. (2009). Colorectal cancer epidemiology: incidence, mortality, survival, and risk factors. *Clin. Colon Rectal Surg.* 22, 191–197.
- Hall, J.A., Cannons, J.L., Grainger, J.R., Dos Santos, L.M., Hand, T.W., Naik, S., Wohlfert, E.A., Chou, D.B., Oldenhove, G., Robinson, M., et al. (2011a). Essential role for retinoic acid in the promotion of CD4(+) T cell effector responses via retinoic acid receptor alpha. *Immunity* 34, 435–447.
- Hall, J.A., Grainger, J.R., Spencer, S.P., and Belkaid, Y. (2011b). The role of retinoic acid in tolerance and immunity. *Immunity* 35, 13–22.
- Hu, B., Elinav, E., Huber, S., Strowig, T., Hao, L., Hafemann, A., Jin, C., Wunderlich, C., Wunderlich, T., Eisenbarth, S.C., and Flavell, R.A. (2013). Microbiota-induced activation of epithelial IL-6 signaling links inflammation-driven inflammation with transmissible cancer. *Proc. Natl. Acad. Sci. USA* 110, 9862–9867.
- Jansa, P., and Forejt, J. (1996). A novel type of retinoic acid response element in the second intron of the mouse H2Kb gene is activated by the RAR/RXR heterodimer. *Nucleic Acids Res.* 24, 694–701.
- Jiménez-Lara, A.M., Clarke, N., Altucci, L., and Gronemeyer, H. (2004). Retinoic-acid-induced apoptosis in leukemia cells. *Trends Mol. Med.* 10, 508–515.
- Kane, M.A., Folias, A.E., Wang, C., and Napoli, J.L. (2008). Quantitative profiling of endogenous retinoic acid in vivo and in vitro by tandem mass spectrometry. *Anal. Chem.* 80, 1702–1708.
- Kedishvili, N.Y. (2013). Enzymology of retinoic acid biosynthesis and degradation. *J. Lipid Res.* 54, 1744–1760.
- Koelzer, V.H., Lugli, A., Dawson, H., Hädrich, M., Berger, M.D., Borner, M., Mallaev, M., Galván, J.A., Amsler, J., Schnüriger, B., et al. (2014). CD8/CD45RO T-cell infiltration in endoscopic biopsies of colorectal cancer predicts nodal metastasis and survival. *J. Transl. Med.* 12, 81.
- Kropotova, E.S., Zinovieva, O.L., Zyryanova, A.F., Dybovaya, V.I., Prasolov, V.S., Beresten, S.F., Oparina, N.Y., and Mashkova, T.D. (2014). Altered expression of multiple genes involved in retinoic acid biosynthesis in human colorectal cancer. *Pathol. Oncol. Res.* 20, 707–717.
- Lee, M.O., Han, S.Y., Jiang, S., Park, J.H., and Kim, S.J. (2000). Differential effects of retinoic acid on growth and apoptosis in human colon cancer cell lines associated with the induction of retinoic acid receptor beta. *Biochem. Pharmacol.* 59, 485–496.
- Marisa, L., de Reyniès, A., Duval, A., Selves, J., Gaub, M.P., Vescovo, L., Etienne-Gralmaud, M.C., Schiappa, R., Guenot, D., Ayadi, M., et al. (2013). Gene expression classification of colon cancer into molecular subtypes: characterization, validation, and prognostic value. *PLoS Med.* 10, e1001453.
- Mark, M., Ghyselinck, N.B., and Chambon, P. (2006). Function of retinoid nuclear receptors: lessons from genetic and pharmacological dissections of the retinoic acid signaling pathway during mouse embryogenesis. *Annu. Rev. Pharmacol. Toxicol.* 46, 451–480.
- Mielke, L.A., Jones, S.A., Raverdeau, M., Higgs, R., Stefanska, A., Groom, J.R., Misiak, A., Dungan, L.S., Sutton, C.E., Streubel, G., et al. (2013). Retinoic acid expression associates with enhanced IL-22 production by $\gamma\delta$ T cells and innate lymphoid cells and attenuation of intestinal inflammation. *J. Exp. Med.* 210, 1117–1124.
- Mucida, D., Park, Y., Kim, G., Turovskaya, O., Scott, I., Kronenberg, M., and Cheroutre, H. (2007). Reciprocal TH17 and regulatory T cell differentiation mediated by retinoic acid. *Science* 317, 256–260.
- Mucida, D., Husain, M.M., Muroi, S., van Wijk, F., Shinnakasu, R., Naoe, Y., Reis, B.S., Huang, Y., Lambolez, F., Docherty, M., et al. (2013). Transcriptional reprogramming of mature CD4⁺ helper T cells generates distinct MHC class II-restricted cytotoxic T lymphocytes. *Nat. Immunol.* 14, 281–289.

- Nagata, T., Segars, J.H., Levi, B.Z., and Ozato, K. (1992). Retinoic acid-dependent transactivation of major histocompatibility complex class I promoters by the nuclear hormone receptor H-2RIIBP in undifferentiated embryonal carcinoma cells. *Proc. Natl. Acad. Sci. USA* *89*, 937–941.
- Nakayama, F., Semba, S., Usami, Y., Chiba, H., Sawada, N., and Yokozaki, H. (2008). Hypermethylation-modulated downregulation of claudin-7 expression promotes the progression of colorectal carcinoma. *Pathobiology* *75*, 177–185.
- Obrochta, K.M., Krois, C.R., Campos, B., and Napoli, J.L. (2015). Insulin regulates retinol dehydrogenase expression and all-trans-retinoic acid biosynthesis through FoxO1. *J. Biol. Chem.* *290*, 7259–7268.
- Reis, B.S., Hoytema van Konijnenburg, D.P., Grivennikov, S.I., and Mucida, D. (2014). Transcription factor T-bet regulates intraepithelial lymphocyte functional maturation. *Immunity* *41*, 244–256.
- Simpson, J.A., Al-Attar, A., Watson, N.F., Scholefield, J.H., Ilyas, M., and Durrant, L.G. (2010). Intratumoral T cell infiltration, MHC class I and STAT1 as biomarkers of good prognosis in colorectal cancer. *Gut* *59*, 926–933.
- Soler, A.P., Miller, R.D., Laughlin, K.V., Carp, N.Z., Klurfeld, D.M., and Mullin, J.M. (1999). Increased tight junctional permeability is associated with the development of colon cancer. *Carcinogenesis* *20*, 1425–1431.
- Tanaka, T., Kohno, H., Suzuki, R., Yamada, Y., Sugie, S., and Mori, H. (2003). A novel inflammation-related mouse colon carcinogenesis model induced by azoxymethane and dextran sodium sulfate. *Cancer Sci.* *94*, 965–973.
- Ullman, T.A., and Itzkowitz, S.H. (2011). Intestinal inflammation and cancer. *Gastroenterology* *140*, 1807–1816.
- Van Wauwe, J.P., Coene, M.C., Goossens, J., Cools, W., and Monbaliu, J. (1990). Effects of cytochrome P-450 inhibitors on the in vivo metabolism of all-trans-retinoic acid in rats. *J. Pharmacol. Exp. Ther.* *252*, 365–369.
- Wirtz, S., Neufert, C., Weigmann, B., and Neurath, M.F. (2007). Chemically induced mouse models of intestinal inflammation. *Nat. Protoc.* *2*, 541–546.
- Zackular, J.P., Baxter, N.T., Iverson, K.D., Sadler, W.D., Petrosino, J.F., Chen, G.Y., and Schloss, P.D. (2013). The gut microbiome modulates colon tumorigenesis. *MBio* *4*, e00692–e13.